



MMWRTM

Morbidity and Mortality Weekly Report

Weekly

December 1, 2006 / Vol. 55 / No. 47

World AIDS Day — December 1, 2006

December 1 marks the 19th observance of World AIDS Day. The theme for this year is “Stop AIDS. Keep the Promise.”

At the end of 2003, an estimated 1.0–1.2 million persons in the United States were living with human immunodeficiency virus (HIV) infection (1). Of these, an estimated 25% were unaware of their infection, underscoring a critical need to expand HIV testing (1).

To address this need, CDC has released revised recommendations for HIV testing (2). These recommendations aim to make HIV testing a routine part of medical care and to further improve rates of HIV diagnosis among pregnant women. Earlier diagnosis of HIV infection will enable more persons to receive life-saving treatment, resulting in improved health and extended life. In addition, the majority of persons who learn they have HIV infection adopt safer behaviors, thereby reducing HIV transmission to others (3). Finally, making HIV testing a routine part of medical care might help reduce the stigma that some associate with an HIV test.

Additional information is available at <http://www.worldaidscampaign.info> and at <http://worldaidsday2006.org>. Surveillance data on HIV/AIDS for 2005 will be available at <http://www.cdc.gov/hiv/topics/surveillance/resources/reports/index.htm#surveillance> (4).

References

1. Glynn M, Rhodes P. Estimated HIV prevalence in the United States at the end of 2003 [Abstract T1-B1101]. Presented at the 2005 National HIV Prevention Conference, Atlanta, GA; June 12–15, 2005.
2. CDC. Revised recommendations for HIV testing of adults, adolescents, and pregnant women in health-care settings. *MMWR* 2006;55(No. RR-14):1–17.
3. Marks G, Crepaz N, Senterfitt JW, Janssen RS. Meta-analysis of high-risk sexual behavior in persons aware and unaware they are infected with HIV in the United States: implications for HIV prevention programs. *J Acquir Immune Defic Syndr* 2005;39:446–53.
4. CDC. HIV/AIDS surveillance report, 2005. Vol. 17. Atlanta, GA: US Department of Health and Human Services, CDC; 2006. In press.

Missed Opportunities for Earlier Diagnosis of HIV Infection — South Carolina, 1997–2005

In September 2006, CDC published revised recommendations for human immunodeficiency virus (HIV) testing in health-care settings to 1) increase early detection of HIV infection by expanding HIV screening of patients and 2) improve access to HIV care and prevention services (e.g., by conducting screening in locations such as emergency departments and urgent-care facilities, where persons who do not otherwise access HIV testing seek health-care services) (1). HIV screening is now recommended for patients aged 13–64 years in all health-care settings after patients are notified that testing will be performed unless they decline (opt-out screening). This represents a substantial change from earlier recommendations to 1) offer HIV testing routinely to all patients only in health-care settings with high HIV prevalence and 2) conduct targeted screening on the basis of risk behaviors for patients in low-prevalence settings (2). This report examines HIV and acquired immunodeficiency syndrome (AIDS) case reporting in South Carolina before the 2006 recommendations were published. During 2001–2005, a total of 4,315 cases of HIV infection were reported in South Carolina. Of these, 41% were in persons (referred to as late testers) in whom AIDS was diagnosed within 1 year of their initial HIV diagnosis* (4).

*The average latent period from HIV infection to onset of AIDS is approximately 10 years (3).

INSIDE

- 1272 Injuries from Motor-Vehicle Collisions with Moose — Maine, 2000–2004
- 1275 Racial/Ethnic Differences Among Youths in Cigarette Smoking and Susceptibility to Start Smoking — United States, 2002–2004
- 1277 Respiratory Syncytial Virus Activity — United States, 2005–2006
- 1279 Notice to Readers

The *MMWR* series of publications is published by the Coordinating Center for Health Information and Service, Centers for Disease Control and Prevention (CDC), U.S. Department of Health and Human Services, Atlanta, GA 30333.

Suggested Citation: Centers for Disease Control and Prevention. [Article title]. *MMWR* 2006;55:[inclusive page numbers].

Centers for Disease Control and Prevention

Julie L. Gerberding, MD, MPH
Director

Tanja Popovic, MD, PhD
(Acting) Chief Science Officer

James W. Stephens, PhD
(Acting) Associate Director for Science

Steven L. Solomon, MD
Director, Coordinating Center for Health Information and Service

Jay M. Bernhardt, PhD, MPH
Director, National Center for Health Marketing

Judith R. Aguilar
(Acting) Director, Division of Health Information Dissemination (Proposed)

Editorial and Production Staff

John S. Moran, MD, MPH
(Acting) Editor, MMWR Series

Suzanne M. Hewitt, MPA
Managing Editor, MMWR Series

Douglas W. Weatherwax
(Acting) Lead Technical Writer-Editor

Catherine H. Bricker, MS
Jude C. Rutledge
Writers-Editors

Beverly J. Holland
Lead Visual Information Specialist

Lynda G. Cupell
Malbea A. LaPete
Visual Information Specialists

Quang M. Doan, MBA
Erica R. Shaver
Information Technology Specialists

Editorial Board

William L. Roper, MD, MPH, Chapel Hill, NC, Chairman
Virginia A. Caine, MD, Indianapolis, IN
David W. Fleming, MD, Seattle, WA
William E. Halperin, MD, DrPH, MPH, Newark, NJ
Margaret A. Hamburg, MD, Washington, DC
King K. Holmes, MD, PhD, Seattle, WA
Deborah Holtzman, PhD, Atlanta, GA
John K. Iglehart, Bethesda, MD
Dennis G. Maki, MD, Madison, WI
Sue Mallonee, MPH, Oklahoma City, OK
Stanley A. Plotkin, MD, Doylestown, PA
Patricia Quinlisk, MD, MPH, Des Moines, IA
Patrick L. Remington, MD, MPH, Madison, WI
Barbara K. Rimer, DrPH, Chapel Hill, NC
John V. Rullan, MD, MPH, San Juan, PR
Anne Schuchat, MD, Atlanta, GA
Dixie E. Snider, MD, MPH, Atlanta, GA
John W. Ward, MD, Atlanta, GA

Of these late testers, 73% made a total of 7,988 visits to a South Carolina health-care facility during 1997–2005 before their first reported positive HIV test. The diagnoses reported for 79% of these visits were not likely to prompt HIV testing under a risk-based testing strategy. These findings suggest that routine, opt-out HIV screening of all patients in health-care settings, rather than risk-based HIV testing, might result in substantially earlier HIV diagnoses in South Carolina.

HIV/AIDS cases have been reportable by patient name in South Carolina since 1986. This analysis used data from the South Carolina HIV/AIDS Reporting System (HARS) for 2001–2005 and included date of first HIV-positive test, date of AIDS diagnosis, and state of residence. Data quality from HARS exceeds CDC minimum standards on reporting timeliness (95% of cases reported within 6 months of a diagnosis) and completeness of reporting (98%, based on a comparison with other data sources) (South Carolina Department of Health and Environment Control [DHEC], unpublished data, 2005).

Since 1996, state law has required that the Office of Research and Statistics (ORS), South Carolina Budget and Control Board receive reports on all diagnoses (classified by *International Classification of Diseases* [ICD] codes) from all emergency departments, hospital inpatient facilities, ambulatory-care facilities, and outpatient surgery facilities within the state. The health-care data for this report were supplied by 60 emergency departments, 62 inpatient facilities, 63 ambulatory-care facilities or outpatient surgery facilities, and 19 free medical clinics in the state, and represent visits that occurred during 1997–2005. ICD diagnoses were grouped into two categories: 1) diagnoses not suggestive of HIV infection and unlikely to have prompted an HIV test (e.g., hypertension, diabetes, and constipation) and 2) diagnoses suggestive of HIV infection that should have prompted an HIV test (e.g., sexually transmitted diseases, symptoms suggestive of acute retroviral syndrome [5], intravenous drug use, and diseases possibly or probably related to HIV infection [6]).

Data from HARS and ORS were linked using several identifiers, including patient name, date of birth, sex, race/ethnicity, and county of residence. This use of the data was approved by DHEC and the ORS Data Oversight Committee. The data were matched in a secured location by authorized persons who were trained in HARS security and confidentiality guidelines. All identifiers were removed from the analysis dataset provided to investigators, who also signed confidentiality agreements.

During 2001–2005, a total of 4,315 persons with HIV infection in South Carolina were reported to HARS, of whom 1,784 (41.3%) were late testers, including 710 (16.5%) who had AIDS diagnosed within 30 days of their initial HIV diagnoses. Women were less likely than men to be late testers; other demographic and risk characteristics of late testers were

similar to those of persons reported to HARS who did not have onset of AIDS within 1 year of their HIV diagnoses. Of the 1,784 late testers, 1,302 (73.0%) had at least one documented visit to a South Carolina health-care facility during 1997–2005 and before the reported date of HIV diagnosis (Table 1).

A total of 7,988 health-care visits were recorded for the 1,302 late testers who had previously visited a health-care facility. Information on transmission category indicated that 441 (33.9%) of these 1,302 persons were identified as injection-drug users or men who have sex with men, persons with high-risk practices that should have prompted HIV screening if risk histories had been elicited during the health-care visits. However, diagnoses reported for 6,277 (78.6%) of these visits were not likely to prompt an HIV test (Table 2). Of the 7,988 visits, 6,303 (78.9%) were to emergency departments, 982 (12.3%) to inpatient settings, 594 (7.4%) to outpatient facilities, and 109 (1.4%) to free clinics. The median time between the visit to a health-care facility and the date of HIV diagnosis was 2.5 years (range: 0–9 years). The 1,302 late testers made a median of four health-care visits before HIV diagnosis (range: 1–132 visits); 280 (21.5%) late testers made only one health-care visit before HIV diagnosis, 567 (43.5%) made two to five previous visits, 259 (19.9%) made six to 10 visits, and 196 (15.1%) made more than 10 visits. Visits occurring

TABLE 1. Number and percentage of HIV-infected persons* with AIDS subsequently diagnosed within 1 year of HIV diagnosis who had visited a health-care facility before date of HIV diagnosis, by selected characteristics — South Carolina, 2001–2005

| Characteristic | No. | (%) |
|-----------------------------------|-------|--------|
| Sex | | |
| Male | 888 | (68.2) |
| Female | 414 | (31.8) |
| Race/Ethnicity† | | |
| Black, non-Hispanic | 1,057 | (81.2) |
| White, non-Hispanic | 214 | (16.4) |
| Hispanic | 21 | (1.6) |
| Age at HIV diagnosis (yrs) | | |
| 13–19 | 23 | (1.8) |
| 20–29 | 202 | (15.5) |
| 30–39 | 430 | (33.0) |
| 40–49 | 411 | (31.6) |
| ≥50 | 236 | (18.1) |
| Transmission category‡ | | |
| Heterosexual | 466 | (35.8) |
| Men who have sex with men (MSM) | 340 | (26.1) |
| Injection-drug user (IDU) | 83 | (6.4) |
| MSM/IDU | 18 | (1.4) |
| Risk not specified | 387 | (29.7) |

* N = 1,302. Reported in South Carolina during 2001–2005.

† Asians/Pacific Islanders, American Indians/Alaska Natives, and persons of multiple races were excluded because numbers were too small for meaningful analysis.

‡ Transfusion recipients and persons with hemophilia were excluded because numbers were too small for meaningful analysis.

TABLE 2. Number and percentage of health-care visits by HIV-infected persons* with AIDS subsequently diagnosed within 1 year of HIV diagnosis who had visited a health-care facility before date of HIV diagnosis, by reported diagnosis — South Carolina, 1997–2005

| Reported diagnosis | No. | (%) |
|---|--------------|----------------|
| Visits with diagnoses likely to prompt an HIV test | | |
| Sexually transmitted disease and related diagnoses | 1,711 | (21.4) |
| Symptoms suggestive of acute retroviral syndrome† | 1,191 | (14.9) |
| Diseases possibly related to HIV‡ | 478 | (6.0) |
| Diseases probably related to HIV¶ | 94 | (1.2) |
| Intravenous drug use and related behaviors | 85 | (1.1) |
| Visits with diagnoses not likely to prompt an HIV test | 6,277 | (78.6) |
| Total visits | 7,988 | (100.0) |

* N = 1,302. Reported in South Carolina during 2001–2005.

† Including fever, lymphadenopathy, and rash.

‡ Including peripheral neuropathy, pneumonia, and thrombocytopenia.

¶ Including cerebral toxoplasmosis, pulmonary tuberculosis, and thrush.

≤6 months before HIV diagnosis accounted for 1,202 (15.1%) of the 7,988 visits; 818 (10.2%) of visits were made >6 months to 1 year before, 1,340 (16.8%) were >1 to 2 years before, 1,337 (16.7%) were >2 to 3 years before, and 3,291 (41.2%) were >3 years before HIV diagnosis.

Reported by: *W Duffus, MD, L Kettinger, MPH, T Stephens, MSPH, J Gibson, MD, South Carolina Dept of Health and Environmental Control; K Weis, MPH, School of Public Health, Univ of South Carolina; M Tyrell, PhD, D Patterson, PhD, C Finney, MPH, WP Bailey, MPH, Office of Research and Statistics, South Carolina Budget and Control Board. B Branson, MD, L Gardner, PhD, PH Kilmarx, MD, Div of HIV/AIDS Prevention, National Center for HIV, Viral Hepatitis, STDs, and Tuberculosis Prevention (proposed), CDC.*

Editorial Note: The findings in this report indicate that HIV-testing practices in South Carolina failed to identify a substantial proportion of HIV-infected persons early in the course of their infection. Early diagnosis of HIV infection is beneficial to the health of the patient (7) and might have a role in limiting further HIV transmission (8). Among the persons identified in this report as late testers (i.e., persons who received an AIDS diagnosis within 1 year of HIV diagnosis), approximately three fourths had visited a South Carolina health-care facility before having HIV diagnosed. Most of the late testers made multiple visits, and most of their visits occurred 1 year or more before diagnosis of HIV infection. These health-care encounters represent missed opportunities for earlier HIV diagnosis. The majority of diagnoses for these previous visits probably would not have prompted HIV testing under a risk-based testing strategy. In addition, the information on transmission category indicated that 441 (33.9%) of 1,302 persons were identified as injection-drug users or men who have sex with men, persons with high-risk practices that should have prompted HIV screening. Combined, these

findings support the new recommendations for routine, opt-out HIV screening of patients in all health-care settings.

In 2004, South Carolina ranked tenth in rate of annual reported AIDS cases in the United States, with 18.1 AIDS cases per 100,000 population (9). The state's data on persons with newly diagnosed HIV in 2004–2005 indicate that a substantial proportion had low CD4+ T cell counts, which would have qualified them for antiretroviral treatment; nearly one third had ≤ 200 cells per mm^3 , and approximately half had ≤ 350 cells per mm^3 (DHEC, unpublished data, 2006). These data also suggest a high prevalence and long duration of undiagnosed HIV infections in South Carolina.

The findings in this report are subject to at least five limitations. First, although HARS and ORS data are comprehensive, certain HIV/AIDS diagnoses and health-care visits probably were not reported. Second, although several variables were available for linking records between the two datasets, matching might not have been successful in all cases. Third, certain late testers might not have been HIV infected at the time of the previous health-care encounters, some of which occurred up to 8 years before AIDS was diagnosed; therefore, those instances might not have been missed opportunities for HIV diagnosis. However, given the long average latent period of approximately 10 years after HIV infection before the onset of AIDS (3), most persons who had AIDS during 2001–2005 would already have been HIV infected during most of their health-care visits beginning in 1997. Fourth, HIV testing might have been recommended but rejected by certain patients during earlier visits; refusal to test might have been related to the stigma that can be associated with risk-based HIV testing. Finally, referral for HIV testing might have occurred during some of the health-care encounters before HIV was diagnosed, so these visits might not represent missed opportunities.

Given the substantial number of health-care encounters in South Carolina during which an earlier diagnosis of HIV might have been made and the high proportion of these visits that would not have suggested the benefit of an HIV test under the risk-based HIV-testing strategy, these findings underscore the need for routine HIV screening of adults and adolescents visiting health-care facilities. The capacity of treatment and preventive services will need to be increased if HIV testing is made routine. Efforts are ongoing in South Carolina to expand these services. The benefit of routine HIV screening, early diagnosis of HIV infection, and linkage of infected persons to these services might be considerable because previous practices of testing based on risk factors or symptoms did not identify a substantial proportion of HIV-infected persons until late in the course of their disease.

References

1. CDC. Revised recommendations for HIV testing of adults, adolescents, and pregnant women in health-care settings. *MMWR* 2006; 55(No. RR-14):1–17.
2. CDC. Revised guidelines for HIV counseling, testing, and referral. *MMWR* 2001;50(No. RR-19).
3. Fauci AS, Pantaleo G, Stanley S, Weissman D. Immunopathogenic mechanisms of HIV infection. *Ann Intern Med* 1996;124:654–63.
4. CDC. Late versus early testing of HIV—16 sites, United States, 2000–2003. *MMWR* 2003;52:581–6.
5. Kahn JO, Walker BD. Acute human immunodeficiency virus type 1 infection. *N Engl J Med* 1998;339:33–9.
6. Greenwald JL, Rich CA, Bessega S, et al. Evaluation of the Centers for Disease Control and Prevention's recommendations regarding routine testing for human immunodeficiency virus by an inpatient service: who are we missing? *Mayo Clin Proc* 2006;81:452–8.
7. Palella FJ, Deloria-Knoll M, Chmiel JS, et al. Survival benefit of initiating antiretroviral therapy in HIV-infected persons in different CD4+ cell strata. *Ann Intern Med* 2003;138:620–6.
8. Marks G, Crepaz N, Senterfitt JW, Janssen RS. Meta-analysis of high-risk sexual behavior in persons aware and unaware they are infected with HIV in the United States: implications for HIV prevention programs. *J Acquir Immune Defic Syndr* 2005;39:446–53.
9. CDC. HIV/AIDS surveillance report. Vol. 16. Atlanta, GA: US Department of Health and Human Services, CDC; 2004:29–30. Available at <http://www.cdc.gov/hiv/topics/surveillance/resources/reports/2004report/pdf/2004surveillancereport.pdf>.

Injuries from Motor-Vehicle Collisions with Moose — Maine, 2000–2004

Moose are among the largest mammals in North America. Standing up to 7.5 feet at the shoulder and weighing up to 1,600 lbs, they are the largest members of the deer family (1–3). Maine's moose population (approximately 29,000) is the biggest in the United States outside of Alaska (4). During a collision with a motor vehicle, a moose usually is struck in the legs, causing its body to roll onto the hood of the vehicle, often collapsing the windshield and roof. As a result, motor-vehicle collisions involving moose are capable of causing substantial injury to vehicle occupants (3). To assess motor-vehicle collisions with moose in Maine and evaluate risk factors for injuries from these types of collisions, the Maine Department of Health and Human Services studied collision reports from 2000–2004. The results of that study indicated that collision rates varied by county but had clear patterns by season and time of day. Variables associated with risk for injury were posted speed limit, type of vehicle, and sex and age of the driver. Measures to reduce collisions with moose should focus on improving driver education programs and developing better engineering controls (e.g., removing roadside vegetation to improve visibility for drivers). In addition, herd management (i.e., decreasing moose population size through hunting) is

currently being used in areas of Maine with high numbers of collisions, although studies are needed to assess its effectiveness.

Information was obtained from motor-vehicle collision reports submitted to the Maine Department of Transportation (DOT) by state, county, and local police during 2000–2004 using a standard form. DOT then entered the report information into two separate data sets: one containing collision information and the other containing driver information. DOT classified collisions into three categories: 1) collisions causing fatal injuries, 2) collisions causing nonfatal injuries, and 3) collisions causing no injuries (5). A nonfatal injury was subcategorized as an incapacitating injury, a nonincapacitating injury, or a possible injury. A noninjury collision was one that resulted in property damage only. Collision rates were calculated using population figures from the 2000 U.S. census. Relative risks (RRs) were calculated for selected exposure variables. Significant ($p < 0.05$) variables were then assessed by logistic regression analysis.

During the 5-year period, 22,516 motor-vehicle collisions with animals were reported in Maine. Of these collisions, 18,289 (81%) were with deer, 3,400 (15%) with moose, and 827 (4%) with other animals. A total of 1,600 injuries (1,583 nonfatal and 17 fatal) were caused by these collisions. Although collisions with moose accounted for only 15% of collisions with animals, they accounted for 803 (50%) of the 1,600 total injuries: 14 (82%) of the 17 fatal injuries and 789 (50%) of the 1,583 nonfatal injuries.

The yearly collision rate with moose was 53 per 100,000 persons overall and ranged from seven to 310 in Maine's 16 counties. Rates were highest in the less populous northern part of the state and lowest in the more populous southeastern part of the state. The majority (2,683 [79%]) of collisions with moose occurred during May–October, with the greatest number of crashes (716 [21%]) occurring in June (Figure 1). The peak time of day for collisions was 10–11 p.m., with 600 collisions (18%); a total of 2,645 (78%) collisions occurred during 6 p.m.–6 a.m. (Figure 2). Occupants of vehicles involved in motor-vehicle collisions with moose were more likely to be injured from the collision when the posted speed limit was ≥ 40 m.p.h. (RR = 1.9, 95% confidence interval [CI] = 1.1–3.3). Neither daylight nor wet road conditions caused by precipitation were significantly associated with a higher risk of being injured in a moose collision. Data regarding locations of collisions were limited to the county level, so particularly high-risk roads or locations could not be identified.

Of the 3,400 collisions with moose, 33 were multivehicle collisions; a total of 3,442 drivers were involved. Because the data assessment did not include identity of the drivers, whether a particular driver had been involved in more than one collision could not be determined. The median age of drivers was

FIGURE 1. Number of motor-vehicle collisions with moose, by month — Maine, 2000–2004





FIGURE 2. Number of motor-vehicle collisions with moose, by time of day — Maine, 2000–2004



43 years (range: 15–90 years). Seventy-three percent of drivers were male, and 99% were considered in normal physical condition at the time the collision occurred; 1% were classified as fatigued, ill, handicapped, or under the influence of alcohol or drugs. Among drivers involved in collisions with moose, drivers of cars had a higher (38%) chance of being injured than drivers of certain other vehicles (10%) (e.g., sport-utility vehicles [SUVs], trucks, vans, buses, farm vehicles, and commercial vehicles) (RR = 3.4, CI = 2.9–3.9).^{*} Drivers aged <25 years were more likely to be injured than drivers aged ≥ 25 years (RR = 1.5, CI = 1.3–1.8), and women were more likely to be injured than men (RR = 1.6, CI = 1.4–1.8).

In logistic regression analysis, only driver age and vehicle type were associated with risk for driver injury. Drivers aged <25 years had higher odds of injury than older drivers (odds ratio [OR] = 1.3, CI = 1.0–1.6). Male drivers of cars had higher odds of injuries (OR = 4.7, CI = 3.7–5.9) than female drivers of cars (OR = 2.8, CI = 1.9–3.9).

^{*} Because few collisions occurred with motorcycles, they were excluded from the analyses but are included in the total numbers of collisions presented in this report.

Reported by: A Pelletier, MD, Div of State and Local Readiness, Coordinating Office for Terrorism Preparedness and Emergency Response, CDC. A Rey, MPH, EIS Officer, CDC.

Editorial Note: Collisions between moose and motor vehicles in Maine cause a disproportionately high number of injuries compared with collisions with other animals. Differences in rates among counties likely are a result of variations in the moose and human population sizes in different areas of the state. The moose population is greater in the northern region of Maine, which has fewer persons than the southern region. The distinct seasonal pattern of collisions with moose (i.e., higher numbers in May–October) correlates with the increased activity of moose during the warmer months and the September–October mating season; in contrast, the deer mating season occurs during October–December, which correlates with higher numbers of deer collisions during these months. The daily time pattern, with higher numbers of collisions occurring during 6 p.m.–6 a.m., seems to correspond with daily patterns of moose activity; moose are more active in the evening and at dawn. In addition, few roads in Maine are lighted, so seeing moose on roads at night is difficult.

The finding that vehicle type was associated with injury in the logistic regression model supports other studies that have found that vehicle type influences likelihood of injury (2). The additional height and mass of larger vehicles such as trucks and SUVs might help protect drivers of these types of vehicles from injury. The association between younger driver age and higher risk for injury might be a result of younger drivers' inexperience and driving habits such as speeding or not using safety belts (6,7). Differences in injury by sex might have been the result of factors that were not included in the logistic regression model (e.g., speed limit, safety-belt use, or driver behavior).

The findings in this report are subject to at least three limitations. First, information on safety-belt use was not included in either data set provided by DOT, although it is recorded in the vehicle collision reports that are submitted to DOT by police. Although the association of safety-belt use with risk for injury could not be assessed in this study, the use of safety belts is the most effective means of reducing fatal and nonfatal injuries in motor-vehicle crashes (8). Second, information regarding the distribution of moose throughout the state was limited. As a result, collision rates based on moose population density could not be calculated. Finally, although the posted speed limit was associated with injury in the bivariate analysis, it was not included in the logistic regression model

because of difficulties associated with merging the collision and driver data sets.

Several public awareness initiatives to prevent motor-vehicle collisions with moose in Maine are ongoing. For example, a statewide campaign involves alerting the public about moose collisions and providing tips for drivers on ways to avoid or decrease the severity of collisions with moose. Brochures are available at libraries, schools, state parks, tourism centers, and other distribution points throughout Maine. In addition, a module on large-animal collisions is a component of Maine Department of Motor Vehicles driver education programs. Other strategies include engineering controls such as clearing roadside vegetation to improve sight lines and placing signs on roads known to have frequent vehicle-moose collisions. Herd management might be an effective strategy in areas with large moose populations. Maine currently manages the size of the moose population through hunting by increasing the number of available moose-hunting permits in areas with high numbers of collisions. Studies are needed to assess the effectiveness of this and other strategies currently being used to reduce the numbers of motor-vehicle collisions with moose.

Acknowledgments

This report is based, in part, on data provided by G Costello, Maine Dept of Transportation, and contributions from A Yartel, MPH, Maine Dept of Health and Human Svcs, J Magri, MD, and A De, PhD, Office of Workforce and Career Development, CDC.

References

1. Bartalucci A, Weinstein B, Dewey T. *Alces alces*. Ann Arbor, MI: University of Michigan, Museum of Zoology; 2000. Available at http://animaldiversity.ummz.umich.edu/site/accounts/information/Alces_alces.html.
2. Farrell TM, Sutton JE, Clark DE, et al. Moose-motor vehicle collisions: an increasing hazard in northern New England. *Arch Surg* 1996;131:377–81.
3. Pynn TP, Pynn BR. Moose and other large animal wildlife vehicle collisions: implications for prevention and emergency care. *J Emerg Nurs* 2004;30:542–7.
4. Seymour T. Southern moose hunt. *Maine Fish Wildl* 2006;47:8–10.
5. Maine Department of Transportation. Collisions between wildlife species and motor vehicles in Maine 1999–2003. Augusta, ME: Maine Department of Transportation; 2004.
6. Boyce TE, Geller ES. An instrumented vehicle assessment of problem behavior and driving style: do younger males really take more risks? *Accid Anal Prev* 2002;34:51–64.
7. McKnight AJ, McKnight AS. Young novice drivers: careless or clueless? *Accid Anal Prevention* 2003;35:921–5.
8. Dinh-Zarr TB, Sleet DA, Shults RA, et al. Reviews of evidence regarding interventions to increase the use of safety belts. *Am J Prev Med* 2001;21(Suppl 4):S48–65.

Racial/Ethnic Differences Among Youths in Cigarette Smoking and Susceptibility to Start Smoking — United States, 2002–2004

Limited information on cigarette smoking in racial/ethnic subpopulations hinders development and implementation of targeted interventions for smoking prevention and cessation. Because of small sample sizes or inadequate study formats, cigarette smoking among youths has been studied mostly in major racial/ethnic populations (e.g., Asian or Hispanic) instead of subsets of these populations (e.g., Vietnamese or Cuban). Data on major population categories might mask differences in tobacco-use prevalence among subpopulations. To assess the prevalence of cigarette smoking among youths aged 12–17 years in six major racial/ethnic populations* and nine Asian or Hispanic subpopulations† in the United States, the Substance Abuse and Mental Health Services Administration and CDC analyzed self-reported data collected during 2002–2004 from the National Survey on Drug Use and Health (NSDUH). This report summarizes the results of that analysis, which indicated that the estimated prevalence of cigarette smoking in this age group ranged from 23.1% for American Indians/Alaska Natives (AI/ANs) to 2.2% for Vietnamese. Implementing tobacco-control programs that include culturally appropriate interventions might help reduce cigarette smoking in racial/ethnic subpopulations.

NSDUH is an annual, in-person household survey that collects information on drug use and abuse from a nationally representative sample of the U.S. civilian, noninstitutionalized population aged ≥ 12 years. The average, weighted, overall response rate for the 2002–2004 surveys was 81% for youths aged 12–17 years, based on a household screening response rate of 91% and an interview response rate of 89%; the final sample size was 68,611. Racial/ethnic classifications by NSDUH were based on standards for classification of federal data (1). Prevalences and 95% confidence intervals (CIs) were calculated; data were weighted to account for different probabilities of selection within strata. Differences in prevalences were considered statistically significant if CIs did not overlap; no other test for statistical significance was performed.

Current cigarette smoking was assessed by asking respondents aged 12–17 years, “During the past 30 days, have you smoked part or all of a cigarette?” Youths who answered “yes” were classified as current smokers. Susceptibility to start smoking

among self-reported nonsmokers was determined by the following two questions: 1) “If one of your best friends offered you a cigarette, would you smoke it?” and 2) “At any time during the next 12 months, do you think that you will smoke a cigarette?” Possible answers were “definitely not,” “probably not,” “probably yes,” and “definitely yes.” Those who answered “definitely not” to both questions were classified as nonsusceptible; those who answered with any other combination of responses were considered susceptible to start smoking.

Among youths, AI/ANs had the greatest cigarette smoking prevalence (23.1%), followed by non-Hispanic whites (14.9%), Hispanics (9.3%), non-Hispanic blacks (6.5%), and Asians (4.3%) (Table 1). Among Asian subpopulations, smoking prevalence ranged from 2.2% for Vietnamese to 6.8% for Koreans; among Hispanic populations, prevalence ranged from 7.3% for Central and South Americans to 11.2% for Cubans. However, none of the differences among Asian subpopulations and Hispanic subpopulations were statistically significant. No significant differences were observed between male and female youths in any of the major populations or subpopulations, except for non-Hispanic white youths, among whom females had a greater prevalence of cigarette smoking (16.0%) than males (13.4%).

A wide range in susceptibility to start smoking was observed among youths who had never smoked (Table 2). Overall, 22.2% were susceptible to start smoking. Youths in the Mexican subpopulation were significantly more susceptible (28.8%) to start smoking than non-Hispanic white (20.8%), non-Hispanic black (23.0%), Cuban (16.4%), Asian Indian (15.4%), Chinese (15.3%), and Vietnamese (13.8%) youths. No significant differences in susceptibility to start smoking were observed between male and female youths in any of the major populations or subpopulations.

Reported by: J Gfroerer, Office of Applied Studies, Substance Abuse and Mental Health Services Admin. R Caraballo, PhD, Office on Smoking and Health, National Center for Chronic Disease Prevention and Health Promotion, CDC.

Editorial Note: The findings in this report indicate that cigarette smoking varied among racial/ethnic subpopulations in addition to major populations of youths aged 12–17 years and that AI/AN youths had the highest prevalence of cigarette smoking in the United States. Differences in smoking prevalence might be attributable to multiple factors, including cigarette prices and discount offers, exposure to antismoking campaigns, and ability to buy cigarettes, all of which can vary by racial/ethnic population (2).

This study also suggests that, overall, approximately one in five nonsmokers aged 12–17 years is susceptible to start smoking. Among the six major populations and nine Asian or

*Major racial/ethnic populations include: Hispanics and the following non-Hispanic populations: white, black or African American, American Indian/Alaska Native, Hawaiian or other Pacific Islander, and Asian.

†Asian subpopulations: Chinese, Filipino, Asian Indian, Korean, and Vietnamese. Hispanic subpopulations: Mexican, Puerto Rican, Central or South American, and Cuban.

TABLE 1. Percentage of youths aged 12–17 years who had smoked one or more cigarettes during the preceding month,* by race/ethnicity and sex — National Survey on Drug Use and Health, United States, 2002–2004

| Race/Ethnicity | Total | | Male | | Female | |
|------------------------------------|-------------|-----------------------|-----------------|--------------------|-------------|--------------------|
| | % | (95% CI) [†] | % | (95% CI) | % | (95% CI) |
| Overall[§] | 12.3 | (12.0–12.7) | 11.8 | (11.4–12.3) | 12.9 | (12.4–13.4) |
| All non-Hispanic [§] | 12.9 | (12.6–13.3) | 12.3 | (11.8–12.8) | 13.6 | (13.1–14.1) |
| White | 14.9 | (14.5–15.4) | 13.9 | (13.4–14.5) | 16.0 | (15.3–16.6) |
| Black or African American | 6.5 | (5.9–7.1) | 7.1 | (6.2–8.1) | 5.9 | (5.1–6.8) |
| American Indian/Alaska Native | 23.1 | (18.9–28.1) | 18.7 | (14.7–23.4) | 28.3 | (21.5–36.3) |
| Hawaiian or other Pacific Islander | 7.1 | (4.0–12.5) | NA [¶] | | 7.8 | (4.0–14.6) |
| Asian [§] | 4.3 | (3.3–5.7) | 5.2 | (3.6–7.4) | 3.4 | (2.2–5.1) |
| Chinese | 2.9 | (1.4–6.0) | 3.7 | (1.4–9.2) | 2.1 | (0.7–6.6) |
| Filipino | 4.6 | (2.3–8.9) | 3.9 | (1.5–9.9) | 5.3 | (2.0–12.9) |
| Asian Indian | 4.5 | (2.4–8.3) | 5.0 | (1.8–12.9) | 4.1 | (2.1–7.7) |
| Korean | 6.8 | (3.3–13.4) | 7.4 | (3.2–16.4) | NA | |
| Vietnamese | 2.2 | (0.7–6.9) | NA | | NA | |
| Hispanic [§] | 9.3 | (8.5–10.1) | 9.2 | (8.1–10.5) | 9.4 | (8.4–10.5) |
| Mexican | 9.0 | (8.0–10.1) | 9.7 | (8.3–11.3) | 8.2 | (7.1–9.6) |
| Puerto Rican | 11.1 | (9.0–13.7) | 9.2 | (6.2–13.4) | 13.4 | (10.3–17.1) |
| Central or South American | 7.3 | (5.4–9.7) | 6.1 | (3.7–9.9) | 8.6 | (6.1–12.1) |
| Cuban | 11.2 | (6.9–17.6) | NA | | 12.2 | (6.9–20.8) |

* As determined by a “yes” response to the question: “During the past 30 days, have you smoked part or all of a cigarette?”

[†] Confidence interval.

[§] Totals include data from respondents reporting other racial/ethnic subpopulations or more than one of those listed.

[¶] Not applicable; values too small for meaningful analysis.

TABLE 2. Percentage of youths aged 12–17 years who had never smoked but were susceptible to start smoking cigarettes,* by race/ethnicity and sex — National Survey on Drug Use and Health, United States, 2002–2004

| Race/Ethnicity | Total | | Male | | Female | |
|------------------------------------|-----------------|-----------------------|-------------|--------------------|-------------|--------------------|
| | % | (95% CI) [†] | % | (95% CI) | % | (95% CI) |
| Overall[§] | 22.2 | (21.8–22.7) | 22.7 | (22.0–23.4) | 21.8 | (21.0–22.5) |
| All non-Hispanic [§] | 21.3 | (20.8–21.8) | 21.8 | (21.1–22.6) | 20.7 | (20.0–21.5) |
| White | 20.8 | (20.3–21.4) | 21.0 | (20.2–21.8) | 20.7 | (19.8–21.5) |
| Black or African American | 23.0 | (21.9–24.2) | 24.1 | (22.5–25.9) | 21.9 | (20.3–23.5) |
| American Indian/Alaska Native | 26.3 | (21.0–32.3) | 32.1 | (24.3–41.0) | 19.4 | (12.7–28.3) |
| Hawaiian or other Pacific Islander | NA [¶] | | NA | | NA | |
| Asian [§] | 18.3 | (15.7–21.2) | 22.1 | (18.1–26.7) | 14.6 | (11.7–18.2) |
| Chinese | 15.3 | (10.4–21.9) | 14.3 | (7.9–24.4) | 16.2 | (10.2–24.7) |
| Filipino | 22.4 | (16.6–29.5) | 26.6 | (17.9–37.5) | 17.9 | (10.8–28.4) |
| Asian Indian | 15.4 | (10.7–21.8) | NA | | 9.9 | (6.0–15.9) |
| Korean | 24.9 | (16.8–35.2) | NA | | NA | |
| Vietnamese | 13.8 | (7.9–23.0) | NA | | NA | |
| Hispanic [§] | 27.0 | (25.6–28.4) | 27.1 | (25.0–29.2) | 26.9 | (24.8–29.2) |
| Mexican | 28.8 | (27.1–30.6) | 29.5 | (27.1–32.0) | 28.1 | (25.5–30.9) |
| Puerto Rican | 23.3 | (19.4–27.7) | 20.2 | (14.7–27.2) | 27.2 | (21.4–33.8) |
| Central or South American | 24.7 | (20.6–29.4) | 25.4 | (19.4–32.5) | 23.9 | (18.7–29.8) |
| Cuban | 16.4 | (11.2–23.4) | 16.6 | (10.4–25.4) | 16.2 | (9.2–26.9) |

* Susceptibility to start smoking among self-reported nonsmokers was determined by the following two questions: 1) “If one of your best friends offered you a cigarette, would you smoke it?” and 2) “At any time during the next 12 months, do you think that you will smoke a cigarette?” Possible answers were “definitely not,” “probably not,” “probably yes,” and “definitely yes.” Those who answered “definitely not” to both questions were classified as nonsusceptible; those who answered with any other combination of responses were considered susceptible to start smoking.

[†] Confidence interval.

[§] Totals include data from respondents reporting other racial/ethnic subpopulations or more than one of those listed.

[¶] Not applicable; values too small for meaningful analysis.

Hispanic subpopulations studied, Mexican youths who had never smoked appeared most susceptible to start smoking. Youths in this subpopulation might need specialized prevention interventions to lower their susceptibility.

Two major public health objectives are 1) to prevent the initiation of cigarette smoking among children, adolescents, and young adults and 2) to help those who already smoke, including children and adolescents, to quit. The overall prevalence of cigarette smoking among high school students declined from 36.4% in 1997 to 23.0% in 2005 (3); however, recent evidence suggests that the reduction in smoking rates over time might have stalled (4).

Children and teens constitute the majority of all new smokers (5). In 2003, cigarette companies spent approximately \$15.2 billion to promote their products, nearly triple their spending in 1996 (6). Conversely, spending by state tobacco-control programs declined from \$749.7 million in 2002 to \$551.0 million in 2006, an amount still less than 3% of the \$21.3 billion that the states received in 2005 from tobacco excise taxes and the 1998 Tobacco Master Settlement Agreement (7). The decline in spending on tobacco-control programs might have been a factor in slowing the progress made in reducing smoking among adolescents (3,8).

The findings in this report are subject to at least four limitations. First, NSDUH surveys are conducted only in English or Spanish, which might have limited participation by some persons (e.g., Asians). Second, the precision of smoking prevalence estimates for certain racial/ethnic subpopulations is low, especially when reported by sex; therefore, differences in prevalence among these subpopulations might not have been detected, and estimates should be interpreted with caution. Third, the data in this report were self-reported in participant households and subject to social-desirability bias (2). However, to reduce this bias, the tobacco-use section in the NSDUH survey was administered using computer-assisted self-interviewing, in which participants read the questions on a computer screen or listened to them through headphones and then entered their responses into the computer. Finally, because of changes in the NSDUH survey methodology in 2002, comparison of the estimates in this report with pre-2002 NSDUH data is not recommended (9).

Sustained, culturally appropriate interventions to prevent youths from starting to smoke or help them to quit might be effective in racial/ethnic populations and subpopulations with high prevalences of cigarette smoking. Effective tobacco-control initiatives might result from comprehensive behavior-based approaches enhanced by 1) using culturally targeted media and education campaigns (10) and 2) increasing the capacities (e.g., for program development) of specific populations to address tobacco use within their communities. To aid

these populations in developing programs, systematic reviews of the effectiveness of interventions to reduce or prevent tobacco use are offered by the *Guide to Community Preventive Services* at <http://www.thecommunityguide.org/tobacco>.

References

- Office of Management and Budget. Standards for the classification of federal data on race and ethnicity. *Federal Register* 1995;60:4674–93.
- CDC. Preventing tobacco use among young people: a report of the Surgeon General. Atlanta, GA: US Department of Health and Human Services, CDC; 1994.
- CDC. Youth risk behavior surveillance—United States, 2005. *MMWR* 2006;55(No. SS-5).
- Johnston LD, O'Malley PM, Bachman JG, Schulenberg JE. Monitoring the future national results on adolescent drug use: overview of key findings, 2005. Bethesda, MD: National Institute on Drug Abuse. In press.
- Substance Abuse and Mental Health Services Administration. Results from the 2005 National Survey on Drug Use and Health: national findings. Rockville, MD: Substance Abuse and Mental Health Services Administration, Office of Applied Studies; 2006.
- Federal Trade Commission. Cigarette report for 2003. Washington, DC: Federal Trade Commission; 2005.
- Campaign for Tobacco-Free Kids. A broken promise to our children: the 1998 state tobacco settlement seven years later. Washington, DC: Campaign for Tobacco Free Kids; 2005. Available at <http://www.tobaccofreekids.org/reports/settlements/2006/fullreport.pdf>.
- Tauras JA, Chaloupka FJ, Farrelly MC, et al. State tobacco control spending and youth smoking. *Am J Public Health* 2005;95:338–4.
- Bowman KR, Chromy JR, Hunter SR, Martin PC, Odom DM, eds. 2003 NSDUH methodological resource book. Rockville, MD: Substance Abuse and Mental Health Services Administration, Office of Applied Studies; 2005.
- Office of Minority Health. Closing the health gap. Washington, DC: US Department of Health and Human Services, Office of Minority Health; 2003. Available at <http://www.omhrc.gov/healthgap>.

Brief Report

Respiratory Syncytial Virus Activity — United States, 2005–2006

Respiratory syncytial virus (RSV) is a major cause of lower respiratory tract infections (LRTIs) (e.g., bronchiolitis and pneumonia) among young children in the United States (1). RSV also causes severe respiratory disease and a substantial number of deaths among older adults (2) and persons with compromised respiratory, cardiac, or immune systems (3). RSV is transmitted person to person through close contact or inhalation of large droplets from a sneeze or cough; infection also can occur through contact with fomites (i.e., contaminated surfaces or objects). In temperate climates, peak RSV activity typically occurs during the winter. This report presents preliminary data on RSV activity reported to the National Respiratory and Enteric Virus Surveillance System (NREVSS) for the weeks ending July 8–November 18, 2006, indicating the onset of the 2006–2007 RSV season, and summarizes RSV trends during July 2005–June 2006. Health-care

providers should consider RSV in the differential diagnosis for persons of all ages with LRTIs and implement appropriate isolation precautions to prevent nosocomial transmission from RSV-infected patients (4). Immune prophylaxis should be considered for certain infants and young children at high risk for complications from RSV infection (e.g., certain premature infants or infants and children with chronic lung and heart disease) (5).

NREVSS is a laboratory-based passive surveillance system that monitors temporal and geographic trends for several respiratory and enteric viruses. The laboratories report weekly to CDC the number of specimens tested for viral pathogens, including RSV, and number of positive test results. During July 2005–June 2006, a total of 71 clinical and public health laboratories in 39 states* and the District of Columbia reported RSV data and are included in this analysis. Eighteen laboratories were excluded because of inconsistent reporting or reporting fewer than 35 weeks of data. A total of 120,503 tests were performed, and 19,533 (16.2%) were positive by antigen-detection testing. National RSV activity† began the week ending November 19, 2005, and continued for 21 weeks until April 1, 2006.

Data were summarized by region (West, East, South, and Central) except those from Florida. Data from Florida came from three laboratories (two in Miami and one in Orlando) and were presented separately because they differed substantially from RSV-detection data from the remainder of the South region (Figure). Regional RSV activity§ was highest during October for Florida, during late December and early January for the South (27 laboratories reporting), during January for the Northeast and Midwest (19 laboratories reporting), and during February for the West (15 laboratories reporting). The Florida RSV season seems similar to those reported from some tropical settings in the Northern Hemisphere (6).

Although 17,736 (91%) RSV detections were reported during November 12, 2005–April 15, 2006, sporadic detections were reported throughout the year. During mid-April through

September 2006, laboratories in 36 states and the District of Columbia reported 1,072 RSV detections; of these, 511 (48%) were from Florida. Additional data from Florida laboratories not participating in NREVSS are available at http://www.doh.state.fl.us/disease_ctrl/epi/RSV/rsv.htm.

For the current reporting period (July 8–November 18, 2006), 62 laboratories in 37 states reported testing for RSV. Preliminary 2006 data suggest that the annual seasonal peak began in Florida during the week ending July 1, in the rest of the South during the week ending October 14, and in the Northeast during the week ending November 11 (Figure).

Health-care providers should consider RSV as a cause of acute respiratory disease in all age groups during the annual seasonal peak. Because the onset of RSV activity can vary among regions and communities, physicians and health-care facilities can consult their local clinical laboratories for the latest data on RSV activity. Although several tests can be used to detect RSV infection in young children, only sensitive reverse transcription–polymerase chain reaction (RT-PCR) assays are sufficient to reliably detect RSV in older children and adults (7). NREVSS expanded reporting to include RT–PCR testing for RSV in 2004. However, these data are not included in the annual summary because of the limited number of laboratories reporting RT–PCR results.

Currently, no vaccine or effective therapy is available for RSV. Infants and children at risk for serious RSV infection can receive immune prophylaxis with monthly doses of a humanized murine anti-RSV monoclonal antibody during the RSV season. Infants and children at risk include those aged <24 months with chronic lung disease who have required medical therapy within 6 months of RSV season onset and those with hemodynamically significant heart disease, and preterm infants born at <32 weeks' gestation or preterm infants born at 32–35 weeks' gestation with at least two additional risk factors (e.g., day care attendance, exposure to environmental pollutants, school-aged siblings, congenital abnormalities of the airways, or neuromuscular disease) during their first RSV season (5). Additional information and updates on RSV national and regional trends are available at <http://www.cdc.gov/ncidod/dvrd/revb/nrevss/index.htm>.

Reported by: *National Respiratory and Enteric Virus Surveillance System collaborating laboratories. AL Fowlkes, AM Fry, MD, LJ Anderson, MD, Div of Viral Diseases, National Center for Immunization and Respiratory Diseases (proposed), CDC.*

References


1. Shay DK, Holman RC, Newman RD, Liu LL, Stout JW, Anderson LJ. Bronchiolitis-associated hospitalizations among U.S. children, 1980–1996. *JAMA* 1999;282:1440–6.
2. Thompson WW, Shay DK, Weintraub E, et al. Mortality associated with influenza and respiratory syncytial virus in the United States. *JAMA* 2003;289:179–86.

* *Northeast:* Connecticut, Massachusetts, New Hampshire, New Jersey, New York, and Rhode Island; *Midwest:* Illinois, Indiana, Minnesota, Missouri, Nebraska, North Dakota, Ohio, South Dakota, and Wisconsin; *South:* Alabama, Arkansas, Delaware, District of Columbia, Georgia, Kentucky, Louisiana, Maryland, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, and Virginia; *West:* Alaska, Arizona, California, Colorado, Hawaii, Montana, Washington, and Wyoming; Florida.

† National RSV activity is defined as the first of 2 consecutive weeks during which 50% of participating laboratories report RSV detections and the mean percentage of specimens positive by antigen detection is >10%.

§ Regional RSV onset and conclusion are defined by NREVSS as the median date that indicates the first of 2 consecutive weeks a participating laboratory reports >10% of specimens testing positive by antigen detection and the last week of >10% positive tests preceding 2 consecutive weeks of <10% positive tests.

FIGURE. Percentage of specimens testing positive for respiratory syncytial virus, by region* and week of report — United States, July 9, 2005–November 18, 2006



* *Northeast*: Connecticut, Massachusetts, New Hampshire, New Jersey, New York, and Rhode Island; *Midwest*: Illinois, Indiana, Minnesota, Missouri, Nebraska, North Dakota, Ohio, South Dakota, and Wisconsin; *South*: Alabama, Arkansas, Delaware, District of Columbia, Georgia, Kentucky, Louisiana, Maryland, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, and Virginia; *West*: Alaska, Arizona, California, Colorado, Hawaii, Montana, Washington, and Wyoming; *Florida*. Data from Florida were presented separately because they differed substantially from RSV-detection data from the remainder of the South region.

- Falsey AR, Hennessey PA, Formica MA, Cox C, Walsh EE. Respiratory syncytial virus infection in elderly and high-risk adults. *N Engl J Med* 2005;352:1749–59.
- CDC. Guidelines for preventing health-care-associated pneumonia, 2003: recommendations of CDC and the Healthcare Infection Control Practices Advisory Committee. *MMWR* 2004;53(No. RR-3).
- Meissner HC, Long SS, American Academy of Pediatrics Committee on Infectious Diseases and Committee on Fetus and Newborn. Revised indications for the use of palivizumab and respiratory syncytial virus immune globulin intravenous for the prevention of respiratory syncytial virus infections. *Pediatrics* 2003;112(6 Pt 1):1447–52.
- Weber MW, Mullholland EK, Greenwood BM. Respiratory syncytial virus infection in tropical and developing countries. *Top Med Int Health* 1998;3:268–80.
- Weinberg GA, Erdman DD, Edwards KM, et al. Superiority of reverse-transcription polymerase chain reaction to conventional viral culture in the diagnosis of acute respiratory tract infections in children. *J Infect Dis* 2004;189:706–10.

Notice to Readers

Epidemiology in Action: Intermediate Analytic Methods Course

CDC and Emory University's Rollins School of Public Health will cosponsor the course *Epidemiology in Action: Intermediate Analytic Methods*, February 26–March 2, 2007, at Emory University, Rollins School of Public Health. The course is designed for practicing public health professionals who have had training and experience in basic applied epidemiology and would like training in additional quantitative skills related to analysis and interpretation of epidemiologic data.

The course includes a review of the fundamentals of descriptive epidemiology and biostatistics, measures of association, normal and binomial distributions, confounding, statistical tests, stratification, logistic regression models, and computer programs as used in epidemiology.

The prerequisite is an introductory course in epidemiology, such as *Epidemiology in Action* or the International Course in Applied Epidemiology. Tuition will be charged. The application deadline is January 26, 2007, or until all slots have been filled.

Additional information and applications are available from Emory University, Hubert Global Health Dept (Attn: Pia), 1518 Clifton Rd. NE, Rm. 746, Atlanta, GA 30322; telephone, (404) 727-3485; fax (404) 727-4590; <http://www.sph.emory.edu/epicourses> or email pvaleri@sph.emory.edu.

Erratum: Vol. 55, No. 46

In the **QuickStats on page 1255**, the third line of the title is missing. The title should read: “Percentage of Persons Aged 22–44 Years at Increased Risk for Human Immunodeficiency Virus (HIV) Infection, by Race/Ethnicity and **Education** — **National Survey of Family Growth,* United States, 2002.**”

TABLE I. Provisional cases of infrequently reported notifiable diseases (<1,000 cases reported during the preceding year) — United States, week ending November 25, 2006 (47th Week)*

| Disease | Current week | Cum 2006 | 5-year weekly average† | Total cases reported for previous years | | | | | States reporting cases during current week (No.) |
|--|--------------|----------|------------------------|---|-------|-------|-------|-------|--|
| | | | | 2005 | 2004 | 2003 | 2002 | 2001 | |
| Anthrax | — | 1 | 0 | — | — | — | 2 | 23 | |
| Botulism: | | | | | | | | | |
| foodborne | — | 13 | 1 | 19 | 16 | 20 | 28 | 39 | |
| infant | 1 | 73 | 2 | 90 | 87 | 76 | 69 | 97 | WA (1) |
| other (wound & unspecified) | — | 43 | 1 | 33 | 30 | 33 | 21 | 19 | |
| Brucellosis | — | 100 | 2 | 122 | 114 | 104 | 125 | 136 | |
| Chancroid | 1 | 27 | 1 | 17 | 30 | 54 | 67 | 38 | MI (1) |
| Cholera | — | 6 | 0 | 8 | 5 | 2 | 2 | 3 | |
| Cyclosporiasis§ | 1 | 108 | 1 | 716 | 171 | 75 | 156 | 147 | OK (1) |
| Diphtheria | — | — | — | — | — | 1 | 1 | 2 | |
| Domestic arboviral diseases§¶: | | | | | | | | | |
| California serogroup | — | 50 | 1 | 80 | 112 | 108 | 164 | 128 | |
| eastern equine | — | 7 | 0 | 21 | 6 | 14 | 10 | 9 | |
| Powassan | — | 1 | — | 1 | 1 | — | 1 | N | |
| St. Louis | — | 7 | 0 | 13 | 12 | 41 | 28 | 79 | |
| western equine | — | — | — | — | — | — | — | — | |
| Ehrlichiosis§: | | | | | | | | | |
| human granulocytic | 3 | 354 | 8 | 790 | 537 | 362 | 511 | 261 | NY (3) |
| human monocytic | 4 | 354 | 5 | 521 | 338 | 321 | 216 | 142 | NY (2), NC (2) |
| human (other & unspecified) | 1 | 160 | 1 | 122 | 59 | 44 | 23 | 6 | NY (1) |
| <i>Haemophilus influenzae</i> **, | | | | | | | | | |
| invasive disease (age <5 yrs): | | | | | | | | | |
| serotype b | — | 9 | 0 | 9 | 19 | 32 | 34 | — | |
| nonserotype b | — | 76 | 3 | 135 | 135 | 117 | 144 | — | |
| unknown serotype | 1 | 172 | 3 | 217 | 177 | 227 | 153 | — | OH (1) |
| Hansen disease§ | — | 66 | 2 | 88 | 105 | 95 | 96 | 79 | |
| Hantavirus pulmonary syndrome§ | — | 30 | 0 | 29 | 24 | 26 | 19 | 8 | |
| Hemolytic uremic syndrome, postdiarrheal§ | — | 219 | 3 | 221 | 200 | 178 | 216 | 202 | |
| Hepatitis C viral, acute | 7 | 673 | 28 | 751 | 713 | 1,102 | 1,835 | 3,976 | PA (2), MI (3), OK (1), WA (1) |
| HIV infection, pediatric (age <13 yrs)§,†† | — | 52 | 5 | 380 | 436 | 504 | 420 | 543 | |
| Influenza-associated pediatric mortality§,§§ | — | 40 | 0 | 45 | — | N | N | N | |
| Listeriosis | 8 | 647 | 13 | 892 | 753 | 696 | 665 | 613 | PA (2), OH (1), MD (2), FL (1), CA (2) |
| Measles¶¶ | — | 44 | 1 | 66 | 37 | 56 | 44 | 116 | |
| Meningococcal disease, invasive***: | | | | | | | | | |
| A, C, Y, & W-135 | 1 | 177 | 4 | 297 | — | — | — | — | WV (1) |
| serogroup B | — | 110 | 3 | 157 | — | — | — | — | |
| other serogroup | 1 | 19 | 0 | 27 | — | — | — | — | IN (1) |
| Mumps | 35 | 6,086 | 5 | 314 | 258 | 231 | 270 | 266 | MN (33), MD (1), CO (1) |
| Plague | — | 16 | 0 | 8 | 3 | 1 | 2 | 2 | |
| Poliomyelitis, paralytic | — | — | — | 1 | — | — | — | — | |
| Psittacosis§ | — | 19 | 0 | 19 | 12 | 12 | 18 | 25 | |
| Q fever§ | — | 136 | 1 | 139 | 70 | 71 | 61 | 26 | |
| Rabies, human | — | 1 | 0 | 2 | 7 | 2 | 3 | 1 | |
| Rubella | — | 9 | — | 11 | 10 | 7 | 18 | 23 | |
| Rubella, congenital syndrome | — | 1 | 0 | 1 | — | 1 | 1 | 3 | |
| SARS-CoV§,††† | — | — | — | — | — | 8 | N | N | |
| Smallpox§ | — | — | — | — | — | — | — | — | |
| Streptococcal toxic-shock syndrome§ | 1 | 85 | 1 | 129 | 132 | 161 | 118 | 77 | OH (1) |
| <i>Streptococcus pneumoniae</i> § | | | | | | | | | |
| invasive disease (age <5 yrs) | 12 | 994 | 18 | 1,257 | 1,162 | 845 | 513 | 498 | RI (2), NY (4), OH (3), MD (2), AZ (1) |
| Syphilis, congenital (age <1 yr) | — | 239 | 8 | 361 | 353 | 413 | 412 | 441 | |
| Tetanus | — | 19 | 1 | 27 | 34 | 20 | 25 | 37 | |
| Toxic-shock syndrome (other than streptococcal)§ | 1 | 89 | 2 | 96 | 95 | 133 | 109 | 127 | CA (1) |
| Trichinellosis | — | 11 | 0 | 19 | 5 | 6 | 14 | 22 | |
| Tularemia§ | — | 80 | 2 | 154 | 134 | 129 | 90 | 129 | |
| Typhoid fever | — | 244 | 5 | 324 | 322 | 356 | 321 | 368 | |
| Vancomycin-intermediate <i>Staphylococcus aureus</i> § | — | 3 | 0 | 2 | — | N | N | N | |
| Vancomycin-resistant <i>Staphylococcus aureus</i> § | — | — | — | 3 | 1 | N | N | N | |
| Yellow fever | — | — | — | — | — | — | 1 | — | |

—: No reported cases. N: Not notifiable. Cum: Cumulative year-to-date counts.

* Incidence data for reporting year 2006 are provisional, whereas data for 2001, 2002, 2003, 2004, and 2005 are finalized.

† Calculated by summing the incidence counts for the current week, the two weeks preceding the current week, and the two weeks following the current week, for a total of 5 preceding years. Additional information is available at <http://www.cdc.gov/epo/dphsi/phs/files/5yearweeklyaverage.pdf>.

§ Not notifiable in all states.

¶ Includes both neuroinvasive and non-neuroinvasive. Updated weekly from reports to the Division of Vector-Borne Infectious Diseases, National Center for Zoonotic, Vector-Borne, and Enteric Diseases (proposed) (ArboNET Surveillance).

** Data for *H. influenzae* (all ages, all serotypes) are available in Table II.

†† Updated monthly from reports to the Division of HIV/AIDS Prevention, National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention (proposed). Implementation of HIV reporting influences the number of cases reported. Pediatric HIV data will not be updated monthly for the remainder of this year due to upgrading of the national HIV/AIDS surveillance data management system. Data for HIV/AIDS are available in Table IV quarterly.

§§ Updated weekly from reports to the Influenza Division, National Center for Immunization and Respiratory Diseases (proposed).

¶¶ No measles cases were reported for the current week.

*** Data for meningococcal disease (all serogroups and unknown serogroups) are available in Table II.

††† Updated weekly from reports to the Division of Viral and Rickettsial Diseases, National Center for Zoonotic, Vector-Borne, and Enteric Diseases (proposed).

TABLE II. Provisional cases of selected notifiable diseases, United States, weeks ending November 25, 2006, and November 26, 2005 (47th Week)*

| Reporting area | Chlamydia† | | | | | Coccidioidomycosis | | | | | Cryptosporidiosis | | | | |
|----------------------|--------------|-------------------|--------|----------|----------|--------------------|-------------------|-------|----------|----------|-------------------|-------------------|-----|----------|----------|
| | Current week | Previous 52 weeks | | Cum 2006 | Cum 2005 | Current week | Previous 52 weeks | | Cum 2006 | Cum 2005 | Current week | Previous 52 weeks | | Cum 2006 | Cum 2005 |
| | | Med | Max | | | | Med | Max | | | | Med | Max | | |
| United States | 6,193 | 19,352 | 35,170 | 860,931 | 863,357 | 78 | 150 | 1,643 | 7,315 | 4,179 | 39 | 73 | 594 | 4,726 | 7,196 |
| New England | 261 | 650 | 1,550 | 29,963 | 28,920 | — | 0 | 0 | — | — | 6 | 3 | 36 | 270 | 337 |
| Connecticut | 54 | 174 | 1,214 | 8,511 | 8,510 | N | 0 | 0 | N | N | — | 0 | 33 | 33 | 77 |
| Maine§ | 41 | 42 | 67 | 2,043 | 2,036 | N | 0 | 0 | N | N | 4 | 0 | 4 | 39 | 30 |
| Massachusetts | 125 | 296 | 607 | 13,930 | 12,923 | — | 0 | 0 | — | — | — | 1 | 14 | 88 | 145 |
| New Hampshire | 4 | 38 | 71 | 1,804 | 1,665 | — | 0 | 0 | — | — | — | 1 | 5 | 47 | 36 |
| Rhode Island | 19 | 63 | 107 | 2,682 | 2,939 | — | 0 | 0 | — | — | — | 0 | 6 | 14 | 13 |
| Vermont§ | 18 | 20 | 43 | 993 | 847 | N | 0 | 0 | N | N | 2 | 0 | 5 | 49 | 36 |
| Mid. Atlantic | 1,107 | 2,410 | 3,696 | 108,583 | 107,077 | — | 0 | 0 | — | — | 10 | 10 | 444 | 523 | 3,129 |
| New Jersey | 63 | 363 | 496 | 16,110 | 17,334 | N | 0 | 0 | N | N | — | 0 | 3 | 11 | 56 |
| New York (Upstate) | 519 | 497 | 1,727 | 21,800 | 21,378 | N | 0 | 0 | N | N | 8 | 3 | 441 | 165 | 2,670 |
| New York City | 314 | 727 | 1,567 | 34,729 | 35,112 | N | 0 | 0 | N | N | — | 2 | 7 | 95 | 141 |
| Pennsylvania | 211 | 768 | 1,104 | 35,944 | 33,253 | N | 0 | 0 | N | N | 2 | 4 | 17 | 252 | 262 |
| E.N. Central | 702 | 3,140 | 12,578 | 141,256 | 146,823 | — | 1 | 3 | 42 | 11 | 2 | 15 | 105 | 1,149 | 1,569 |
| Illinois | 322 | 977 | 1,697 | 47,285 | 45,665 | — | 0 | 0 | — | — | — | 2 | 18 | 140 | 154 |
| Indiana | — | 387 | 478 | 17,243 | 18,119 | N | 0 | 0 | N | N | 1 | 1 | 18 | 90 | 79 |
| Michigan | 256 | 658 | 9,888 | 31,236 | 25,377 | — | 0 | 3 | 36 | 11 | — | 2 | 8 | 129 | 103 |
| Ohio | 46 | 636 | 1,424 | 28,189 | 39,189 | — | 0 | 2 | 6 | — | 1 | 5 | 33 | 335 | 751 |
| Wisconsin | 78 | 385 | 531 | 17,303 | 18,473 | N | 0 | 0 | N | N | — | 5 | 53 | 455 | 482 |
| W.N. Central | 511 | 1,160 | 1,455 | 53,326 | 53,247 | — | 0 | 12 | 1 | 4 | 2 | 12 | 77 | 801 | 584 |
| Iowa | 99 | 159 | 225 | 7,495 | 6,661 | N | 0 | 0 | N | N | — | 1 | 28 | 167 | 120 |
| Kansas | 106 | 150 | 269 | 6,479 | 6,665 | N | 0 | 0 | N | N | — | 1 | 8 | 77 | 36 |
| Minnesota | — | 236 | 347 | 10,091 | 11,120 | — | 0 | 12 | — | 3 | — | 3 | 22 | 214 | 127 |
| Missouri | 139 | 440 | 612 | 20,535 | 20,224 | — | 0 | 1 | 1 | 1 | 2 | 2 | 21 | 174 | 244 |
| Nebraska§ | 107 | 96 | 176 | 4,858 | 4,580 | N | 0 | 0 | N | N | — | 1 | 16 | 88 | 26 |
| North Dakota | 12 | 33 | 61 | 1,500 | 1,511 | N | 0 | 0 | N | N | — | 0 | 4 | 9 | 1 |
| South Dakota | 48 | 51 | 116 | 2,368 | 2,486 | N | 0 | 0 | N | N | — | 1 | 7 | 72 | 30 |
| S. Atlantic | 1,653 | 3,695 | 4,936 | 167,079 | 158,817 | — | 0 | 1 | 3 | 2 | 16 | 15 | 70 | 1,053 | 698 |
| Delaware | 59 | 67 | 92 | 3,212 | 3,068 | N | 0 | 0 | N | N | — | 0 | 3 | 15 | 6 |
| District of Columbia | 38 | 53 | 138 | 2,629 | 3,418 | — | 0 | 0 | — | — | — | 0 | 2 | 14 | 15 |
| Florida | 423 | 964 | 1,157 | 44,022 | 38,646 | N | 0 | 0 | N | N | 6 | 6 | 32 | 504 | 323 |
| Georgia | 21 | 685 | 2,142 | 29,247 | 28,427 | — | 0 | 0 | — | — | 5 | 4 | 12 | 224 | 139 |
| Maryland§ | 205 | 328 | 487 | 15,895 | 16,677 | — | 0 | 1 | 3 | 2 | 1 | 0 | 3 | 19 | 30 |
| North Carolina | 572 | 593 | 1,772 | 30,218 | 28,575 | N | 0 | 0 | N | N | 3 | 1 | 11 | 93 | 84 |
| South Carolina§ | 108 | 347 | 1,452 | 17,622 | 17,075 | N | 0 | 0 | N | N | — | 1 | 13 | 122 | 23 |
| Virginia§ | 207 | 430 | 840 | 21,445 | 20,459 | N | 0 | 0 | N | N | — | 1 | 6 | 52 | 64 |
| West Virginia | 20 | 58 | 227 | 2,789 | 2,472 | N | 0 | 0 | N | N | 1 | 0 | 3 | 10 | 14 |
| E.S. Central | 433 | 1,420 | 1,947 | 66,452 | 62,731 | — | 0 | 0 | — | — | 3 | 3 | 12 | 174 | 210 |
| Alabama§ | 48 | 407 | 756 | 18,717 | 14,922 | N | 0 | 0 | N | N | 3 | 1 | 10 | 80 | 25 |
| Kentucky | 61 | 163 | 613 | 7,876 | 7,772 | N | 0 | 0 | N | N | — | 1 | 5 | 35 | 139 |
| Mississippi | — | 365 | 807 | 16,845 | 19,121 | — | 0 | 0 | — | — | — | 0 | 3 | 16 | 2 |
| Tennessee§ | 324 | 512 | 608 | 23,014 | 20,916 | N | 0 | 0 | N | N | — | 1 | 5 | 43 | 44 |
| W.S. Central | 125 | 2,177 | 3,605 | 97,353 | 99,521 | — | 0 | 1 | 2 | — | — | 4 | 44 | 322 | 220 |
| Arkansas | 77 | 153 | 335 | 7,386 | 7,780 | — | 0 | 1 | 1 | — | — | 0 | 2 | 20 | 6 |
| Louisiana | 48 | 245 | 607 | 11,854 | 15,771 | — | 0 | 1 | 1 | N | — | 0 | 9 | 67 | 81 |
| Oklahoma | — | 227 | 2,159 | 11,232 | 10,463 | N | 0 | 0 | N | N | — | 1 | 4 | 38 | 41 |
| Texas§ | — | 1,458 | 1,903 | 66,881 | 65,507 | N | 0 | 0 | N | N | — | 2 | 35 | 197 | 92 |
| Mountain | 262 | 1,025 | 1,839 | 46,119 | 56,307 | 62 | 109 | 452 | 5,018 | 2,707 | — | 3 | 39 | 359 | 133 |
| Arizona | 232 | 368 | 881 | 17,294 | 18,824 | 62 | 106 | 448 | 4,899 | 2,605 | — | 0 | 3 | 24 | 10 |
| Colorado | — | 145 | 395 | 5,480 | 13,932 | N | 0 | 0 | N | N | — | 1 | 7 | 67 | 49 |
| Idaho§ | — | 48 | 191 | 2,333 | 2,454 | N | 0 | 0 | N | N | — | 0 | 5 | 35 | 14 |
| Montana§ | 23 | 45 | 195 | 2,309 | 2,088 | N | 0 | 0 | N | N | — | 1 | 26 | 131 | 18 |
| Nevada§ | 7 | 85 | 432 | 4,569 | 6,418 | — | 1 | 4 | 52 | 62 | — | 0 | 1 | 9 | 11 |
| New Mexico§ | — | 189 | 339 | 8,477 | 7,442 | — | 0 | 3 | 13 | 19 | — | 0 | 5 | 27 | 17 |
| Utah | — | 94 | 176 | 4,470 | 4,105 | — | 1 | 3 | 52 | 18 | — | 0 | 3 | 18 | 11 |
| Wyoming | — | 27 | 54 | 1,187 | 1,044 | — | 0 | 2 | 2 | 3 | — | 0 | 11 | 48 | 3 |
| Pacific | 1,139 | 3,320 | 5,079 | 150,800 | 149,914 | 16 | 46 | 1,179 | 2,249 | 1,455 | — | 2 | 52 | 75 | 316 |
| Alaska | 32 | 82 | 152 | 3,657 | 3,822 | — | 0 | 0 | — | — | — | 0 | 1 | 4 | 3 |
| California | 653 | 2,570 | 4,231 | 118,552 | 116,258 | 16 | 46 | 1,179 | 2,249 | 1,455 | — | 0 | 14 | — | 189 |
| Hawaii | 2 | 101 | 135 | 4,627 | 4,985 | N | 0 | 0 | N | N | — | 0 | 1 | 4 | 1 |
| Oregon§ | 209 | 165 | 315 | 7,937 | 8,075 | N | 0 | 0 | N | N | — | 1 | 7 | 67 | 67 |
| Washington | 243 | 348 | 604 | 16,027 | 16,774 | N | 0 | 0 | N | N | — | 0 | 38 | — | 56 |
| American Samoa | U | 0 | 46 | U | U | U | 0 | 0 | U | U | U | 0 | 0 | U | U |
| C.N.M.I. | U | 0 | 0 | U | U | U | 0 | 0 | U | U | U | 0 | 0 | U | U |
| Guam | — | 17 | 27 | — | 768 | — | 0 | 0 | — | — | — | 0 | 0 | — | — |
| Puerto Rico | — | 82 | 187 | 3,855 | 3,678 | N | 0 | 0 | N | N | N | 0 | 0 | N | N |
| U.S. Virgin Islands | — | 5 | 16 | 178 | 196 | — | 0 | 0 | — | — | — | 0 | 0 | — | — |

C.N.M.I.: Commonwealth of Northern Mariana Islands.

U: Unavailable. —: No reported cases. N: Not notifiable. Cum: Cumulative year-to-date counts. Med: Median. Max: Maximum.

* Incidence data for reporting year 2006 is provisional.

† Chlamydia refers to genital infections caused by *Chlamydia trachomatis*.

§ Contains data reported through the National Electronic Disease Surveillance System (NEDSS).

TABLE II. (Continued) Provisional cases of selected notifiable diseases, United States, weeks ending November 25, 2006, and November 26, 2005 (47th Week)*

| Reporting area | Giardiasis | | | | | Gonorrhea | | | | | Haemophilus influenzae, invasive All ages, all serotypes | | | | |
|----------------------|--------------|-------------------|-------|----------|----------|--------------|-------------------|--------|----------|----------|---|-------------------|-----|----------|----------|
| | Current week | Previous 52 weeks | | Cum 2006 | Cum 2005 | Current week | Previous 52 weeks | | Cum 2006 | Cum 2005 | Current week | Previous 52 weeks | | Cum 2006 | Cum 2005 |
| | | Med | Max | | | | Med | Max | | | | Med | Max | | |
| United States | 138 | 319 | 1,029 | 15,341 | 17,388 | 1,893 | 6,584 | 14,136 | 299,360 | 297,772 | 19 | 39 | 142 | 1,770 | 1,996 |
| New England | 13 | 23 | 75 | 1,094 | 1,562 | 40 | 110 | 288 | 5,053 | 5,173 | 2 | 2 | 19 | 138 | 150 |
| Connecticut | — | 1 | 37 | 268 | 347 | 14 | 42 | 241 | 2,017 | 2,190 | — | 0 | 9 | 43 | 44 |
| Maine† | 6 | 2 | 13 | 168 | 192 | 2 | 2 | 8 | 117 | 126 | — | 0 | 4 | 19 | 10 |
| Massachusetts | — | 9 | 18 | 357 | 685 | 14 | 47 | 98 | 2,233 | 2,254 | — | 1 | 7 | 52 | 72 |
| New Hampshire | — | 0 | 9 | 27 | 59 | 1 | 3 | 9 | 175 | 160 | — | 0 | 2 | 9 | 8 |
| Rhode Island | 2 | 0 | 25 | 102 | 107 | 7 | 9 | 19 | 450 | 389 | 2 | 0 | 7 | 6 | 7 |
| Vermont† | 5 | 3 | 12 | 172 | 172 | 2 | 1 | 4 | 61 | 54 | — | 0 | 2 | 9 | 9 |
| Mid. Atlantic | 43 | 62 | 254 | 2,987 | 3,134 | 233 | 646 | 1,014 | 29,113 | 30,835 | 4 | 7 | 30 | 339 | 385 |
| New Jersey | — | 9 | 13 | 339 | 418 | 47 | 102 | 160 | 4,580 | 5,136 | — | 0 | 4 | — | 80 |
| New York (Upstate) | 37 | 24 | 227 | 1,141 | 1,093 | 77 | 122 | 455 | 5,619 | 6,290 | 3 | 3 | 27 | 129 | 106 |
| New York City | 1 | 15 | 29 | 782 | 823 | 50 | 175 | 378 | 8,712 | 9,419 | — | 2 | 6 | 78 | 73 |
| Pennsylvania | 5 | 15 | 32 | 725 | 800 | 59 | 224 | 399 | 10,202 | 9,990 | 1 | 3 | 8 | 132 | 126 |
| E.N. Central | 8 | 47 | 82 | 2,205 | 3,059 | 242 | 1,279 | 7,047 | 57,621 | 59,695 | 4 | 5 | 14 | 246 | 337 |
| Illinois | — | 9 | 21 | 359 | 718 | 116 | 378 | 711 | 18,051 | 18,081 | — | 1 | 6 | 47 | 113 |
| Indiana | N | 0 | 0 | N | N | — | 161 | 244 | 7,670 | 7,313 | 1 | 1 | 11 | 73 | 58 |
| Michigan | — | 14 | 37 | 625 | 729 | 64 | 261 | 5,880 | 13,160 | 10,310 | — | 0 | 3 | 20 | 23 |
| Ohio | 8 | 16 | 32 | 744 | 733 | 16 | 303 | 648 | 12,828 | 18,731 | 3 | 2 | 6 | 79 | 103 |
| Wisconsin | — | 10 | 40 | 477 | 879 | 46 | 133 | 172 | 5,912 | 5,260 | — | 0 | 4 | 27 | 40 |
| W.N. Central | 4 | 28 | 260 | 1,610 | 2,056 | 143 | 370 | 444 | 16,893 | 16,919 | 1 | 2 | 15 | 137 | 102 |
| Iowa | — | 5 | 15 | 263 | 257 | 19 | 37 | 62 | 1,665 | 1,472 | — | 0 | 1 | 2 | — |
| Kansas | — | 3 | 11 | 180 | 192 | 28 | 41 | 124 | 1,815 | 2,333 | — | 0 | 3 | 14 | 14 |
| Minnesota | — | 1 | 238 | 481 | 894 | 1 | 63 | 105 | 2,613 | 3,161 | — | 0 | 9 | 72 | 40 |
| Missouri | 3 | 9 | 28 | 492 | 472 | 55 | 190 | 252 | 9,059 | 8,499 | — | 0 | 6 | 32 | 31 |
| Nebraska† | 1 | 2 | 9 | 105 | 111 | 33 | 26 | 56 | 1,284 | 1,038 | — | 0 | 2 | 8 | 14 |
| North Dakota | — | 0 | 7 | 17 | 17 | — | 3 | 7 | 115 | 103 | 1 | 0 | 3 | 9 | 3 |
| South Dakota | — | 1 | 5 | 72 | 113 | 7 | 6 | 15 | 342 | 313 | — | 0 | 0 | — | — |
| S. Atlantic | 22 | 50 | 95 | 2,391 | 2,500 | 668 | 1,607 | 2,334 | 74,948 | 70,214 | 4 | 10 | 24 | 475 | 475 |
| Delaware | — | 1 | 4 | 36 | 53 | 27 | 27 | 44 | 1,336 | 806 | — | 0 | 1 | 1 | — |
| District of Columbia | — | 1 | 4 | 57 | 51 | 43 | 35 | 61 | 1,680 | 1,926 | — | 0 | 2 | 7 | 9 |
| Florida | 13 | 19 | 44 | 1,022 | 880 | 205 | 458 | 549 | 20,780 | 17,916 | 4 | 3 | 9 | 155 | 120 |
| Georgia | 3 | 11 | 28 | 524 | 675 | 13 | 337 | 1,014 | 14,807 | 13,363 | — | 2 | 6 | 89 | 101 |
| Maryland† | 3 | 3 | 11 | 194 | 194 | 77 | 125 | 188 | 5,847 | 6,305 | — | 1 | 5 | 63 | 69 |
| North Carolina | N | 0 | 0 | N | N | 180 | 310 | 766 | 15,680 | 13,804 | — | 0 | 9 | 51 | 72 |
| South Carolina† | — | 1 | 7 | 91 | 101 | 52 | 150 | 704 | 7,977 | 7,937 | — | 1 | 3 | 32 | 32 |
| Virginia† | 3 | 8 | 50 | 435 | 501 | 63 | 130 | 288 | 5,939 | 7,519 | — | 1 | 8 | 58 | 46 |
| West Virginia | — | 0 | 6 | 32 | 45 | 8 | 18 | 43 | 902 | 638 | — | 0 | 4 | 19 | 26 |
| E.S. Central | 2 | 8 | 41 | 480 | 390 | 182 | 568 | 870 | 26,890 | 25,264 | 2 | 2 | 7 | 94 | 107 |
| Alabama† | — | 5 | 29 | 268 | 183 | 18 | 188 | 311 | 8,587 | 8,359 | — | 0 | 5 | 21 | 17 |
| Kentucky | N | 0 | 0 | N | N | 38 | 56 | 180 | 2,866 | 2,735 | — | 0 | 1 | 5 | 12 |
| Mississippi | — | 0 | 0 | — | — | — | 143 | 436 | 6,643 | 6,394 | — | 0 | 1 | 3 | — |
| Tennessee† | 2 | 4 | 12 | 212 | 207 | 126 | 191 | 238 | 8,794 | 7,776 | 2 | 1 | 4 | 65 | 78 |
| W. S. Central | 3 | 5 | 31 | 279 | 303 | 103 | 898 | 1,430 | 42,161 | 40,677 | 1 | 1 | 15 | 61 | 105 |
| Arkansas | — | 2 | 8 | 126 | 78 | 54 | 81 | 142 | 3,850 | 4,068 | — | 0 | 2 | 7 | 7 |
| Louisiana | — | 0 | 5 | 34 | 59 | 49 | 148 | 354 | 7,361 | 8,769 | — | 0 | 3 | 11 | 35 |
| Oklahoma | 3 | 2 | 24 | 119 | 166 | — | 82 | 764 | 4,189 | 4,157 | 1 | 1 | 14 | 43 | 56 |
| Texas† | N | 0 | 0 | N | N | — | 567 | 915 | 26,761 | 23,683 | — | 0 | 1 | — | 7 |
| Mountain | 11 | 30 | 66 | 1,519 | 1,413 | 66 | 222 | 552 | 10,562 | 12,057 | 1 | 4 | 8 | 174 | 199 |
| Arizona | 1 | 3 | 36 | 141 | 134 | 59 | 92 | 201 | 4,286 | 4,345 | — | 1 | 7 | 79 | 98 |
| Colorado | 5 | 9 | 33 | 504 | 496 | — | 45 | 85 | 2,067 | 2,900 | 1 | 1 | 4 | 45 | 39 |
| Idaho† | 5 | 3 | 12 | 172 | 142 | — | 2 | 15 | 139 | 106 | — | 0 | 1 | 6 | 5 |
| Montana† | — | 2 | 11 | 99 | 67 | 5 | 3 | 20 | 178 | 136 | — | 0 | 0 | — | — |
| Nevada† | — | 1 | 8 | 85 | 108 | 2 | 25 | 194 | 1,475 | 2,494 | — | 0 | 1 | 1 | 14 |
| New Mexico† | — | 1 | 6 | 63 | 81 | — | 32 | 65 | 1,540 | 1,366 | — | 0 | 4 | 24 | 25 |
| Utah | — | 7 | 24 | 419 | 359 | — | 18 | 25 | 767 | 637 | — | 0 | 4 | 16 | 9 |
| Wyoming | — | 1 | 4 | 36 | 26 | — | 2 | 6 | 110 | 73 | — | 0 | 1 | 3 | 9 |
| Pacific | 32 | 59 | 202 | 2,776 | 2,971 | 216 | 795 | 967 | 36,119 | 36,938 | — | 2 | 15 | 106 | 136 |
| Alaska | 1 | 1 | 17 | 96 | 105 | 7 | 11 | 24 | 501 | 523 | — | 0 | 2 | 9 | 27 |
| California | 16 | 42 | 105 | 1,972 | 2,110 | 127 | 654 | 834 | 29,775 | 30,742 | — | 0 | 9 | 27 | 56 |
| Hawaii | — | 1 | 3 | 40 | 60 | 4 | 18 | 29 | 794 | 927 | — | 0 | 1 | 16 | 9 |
| Oregon† | 2 | 8 | 14 | 350 | 384 | 28 | 27 | 49 | 1,208 | 1,415 | — | 1 | 6 | 52 | 44 |
| Washington | 13 | 6 | 90 | 318 | 312 | 50 | 76 | 142 | 3,841 | 3,331 | — | 0 | 4 | 2 | — |
| American Samoa | U | 0 | 0 | U | U | U | 0 | 2 | U | U | U | 0 | 0 | U | U |
| C.N.M.I. | U | 0 | 0 | U | U | U | 0 | 0 | U | U | U | 0 | 0 | U | U |
| Guam | — | 0 | 0 | — | 11 | — | 3 | 15 | — | 82 | — | 0 | 1 | — | 14 |
| Puerto Rico | — | 1 | 12 | 77 | 243 | — | 5 | 16 | 239 | 332 | — | 0 | 0 | — | 4 |
| U.S. Virgin Islands | — | 0 | 0 | — | — | — | 0 | 5 | 30 | 45 | — | 0 | 0 | — | — |

C.N.M.I.: Commonwealth of Northern Mariana Islands.

U: Unavailable. —: No reported cases. N: Not notifiable. Cum: Cumulative year-to-date counts. Med: Median. Max: Maximum.

* Incidence data for reporting year 2006 is provisional.

† Contains data reported through the National Electronic Disease Surveillance System (NEDSS).

TABLE II. (Continued) Provisional cases of selected notifiable diseases, United States, weeks ending November 25, 2006, and November 26, 2005 (47th Week)*

| Reporting area | Hepatitis (viral, acute), by type | | | | | | | | | | Legionellosis | | | | |
|----------------------|-----------------------------------|-------------------|-----|----------|----------|--------------|-------------------|-----|----------|----------|---------------|-------------------|-----|----------|----------|
| | A | | | | | B | | | | | | | | | |
| | Current week | Previous 52 weeks | | Cum 2006 | Cum 2005 | Current week | Previous 52 weeks | | Cum 2006 | Cum 2005 | Current week | Previous 52 weeks | | Cum 2006 | Cum 2005 |
| | Med | Max | | | | Med | Max | | | | Med | Max | | | |
| United States | 23 | 68 | 245 | 3,065 | 3,801 | 17 | 84 | 574 | 3,658 | 4,379 | 23 | 41 | 127 | 2,173 | 2,013 |
| New England | 2 | 3 | 20 | 154 | 432 | — | 2 | 8 | 87 | 142 | 2 | 2 | 12 | 115 | 143 |
| Connecticut | — | 1 | 2 | 37 | 48 | — | 0 | 3 | 29 | 44 | 1 | 0 | 9 | 49 | 33 |
| Maine† | — | 0 | 2 | 6 | 4 | — | 0 | 2 | 19 | 12 | — | 0 | 2 | 8 | 7 |
| Massachusetts | — | 0 | 6 | 51 | 279 | — | 0 | 5 | 14 | 48 | — | 0 | 4 | 27 | 64 |
| New Hampshire | — | 0 | 16 | 37 | 80 | — | 0 | 2 | 13 | 29 | — | 0 | 1 | 1 | 9 |
| Rhode Island | 2 | 0 | 4 | 14 | 15 | — | 0 | 4 | 9 | 3 | 1 | 0 | 10 | 22 | 21 |
| Vermont† | — | 0 | 2 | 9 | 6 | — | 0 | 1 | 3 | 6 | — | 0 | 2 | 8 | 9 |
| Mid. Atlantic | 1 | 6 | 17 | 321 | 606 | 2 | 8 | 55 | 382 | 598 | 5 | 15 | 47 | 823 | 697 |
| New Jersey | — | 1 | 6 | 71 | 144 | — | 2 | 8 | 96 | 219 | — | 1 | 11 | 96 | 116 |
| New York (Upstate) | 1 | 1 | 14 | 84 | 92 | 1 | 1 | 43 | 57 | 53 | 2 | 6 | 30 | 305 | 176 |
| New York City | — | 2 | 10 | 107 | 276 | — | 2 | 5 | 80 | 124 | — | 2 | 14 | 124 | 112 |
| Pennsylvania | — | 1 | 5 | 59 | 94 | 1 | 3 | 9 | 149 | 202 | 3 | 4 | 18 | 298 | 293 |
| E.N. Central | — | 6 | 13 | 283 | 338 | 1 | 8 | 24 | 367 | 529 | 4 | 8 | 26 | 427 | 412 |
| Illinois | — | 1 | 4 | 61 | 118 | — | 1 | 7 | 60 | 149 | — | 0 | 4 | 21 | 57 |
| Indiana | — | 0 | 5 | 30 | 19 | — | 0 | 17 | 53 | 40 | — | 0 | 4 | 34 | 30 |
| Michigan | — | 2 | 8 | 106 | 106 | — | 3 | 6 | 129 | 173 | 1 | 2 | 9 | 123 | 109 |
| Ohio | — | 1 | 4 | 49 | 49 | 1 | 2 | 10 | 117 | 123 | 3 | 4 | 19 | 213 | 183 |
| Wisconsin | — | 1 | 4 | 37 | 46 | — | 0 | 2 | 8 | 44 | — | 0 | 5 | 36 | 33 |
| W.N. Central | — | 2 | 30 | 122 | 84 | 1 | 4 | 22 | 151 | 249 | — | 1 | 15 | 74 | 93 |
| Iowa | — | 0 | 2 | 11 | 19 | — | 0 | 3 | 16 | 27 | — | 0 | 3 | 10 | 8 |
| Kansas | — | 0 | 5 | 27 | 16 | 1 | 0 | 2 | 11 | 27 | — | 0 | 2 | 6 | 3 |
| Minnesota | — | 0 | 29 | 16 | 3 | — | 0 | 13 | 23 | 29 | — | 0 | 11 | 24 | 26 |
| Missouri | — | 1 | 3 | 43 | 30 | — | 2 | 6 | 78 | 135 | — | 0 | 3 | 20 | 29 |
| Nebraska† | — | 0 | 2 | 17 | 15 | — | 0 | 3 | 20 | 24 | — | 0 | 2 | 9 | 4 |
| North Dakota | — | 0 | 2 | — | — | — | 0 | 0 | — | — | — | 0 | 1 | — | 2 |
| South Dakota | — | 0 | 3 | 8 | 1 | — | 0 | 1 | 3 | 7 | — | 0 | 1 | 5 | 21 |
| S. Atlantic | 6 | 10 | 29 | 515 | 671 | 5 | 23 | 66 | 1,043 | 1,261 | 4 | 8 | 19 | 398 | 384 |
| Delaware | — | 0 | 2 | 12 | 6 | — | 1 | 4 | 44 | 30 | — | 0 | 2 | 11 | 16 |
| District of Columbia | 1 | 0 | 2 | 8 | 4 | — | 0 | 2 | 7 | 11 | 1 | 0 | 5 | 30 | 12 |
| Florida | 4 | 4 | 13 | 198 | 269 | 3 | 8 | 19 | 378 | 437 | 2 | 3 | 9 | 146 | 104 |
| Georgia | 1 | 1 | 5 | 57 | 117 | 2 | 3 | 8 | 151 | 189 | — | 0 | 4 | 20 | 37 |
| Maryland† | — | 1 | 6 | 61 | 69 | — | 3 | 10 | 138 | 142 | — | 1 | 7 | 83 | 105 |
| North Carolina | — | 0 | 20 | 95 | 82 | — | 0 | 23 | 147 | 150 | 1 | 0 | 5 | 34 | 31 |
| South Carolina† | — | 0 | 3 | 23 | 40 | — | 2 | 7 | 73 | 142 | — | 0 | 1 | 4 | 15 |
| Virginia† | — | 1 | 11 | 55 | 80 | — | 1 | 18 | 56 | 123 | — | 1 | 7 | 57 | 44 |
| West Virginia | — | 0 | 3 | 6 | 4 | — | 0 | 18 | 49 | 37 | — | 0 | 3 | 13 | 20 |
| E.S. Central | 2 | 2 | 8 | 118 | 229 | 2 | 6 | 16 | 315 | 341 | 1 | 1 | 9 | 93 | 81 |
| Alabama† | — | 0 | 3 | 18 | 42 | 2 | 2 | 12 | 110 | 87 | — | 0 | 2 | 10 | 13 |
| Kentucky | — | 0 | 5 | 31 | 24 | — | 1 | 5 | 66 | 66 | — | 0 | 5 | 38 | 29 |
| Mississippi | 1 | 0 | 1 | 9 | 19 | — | 0 | 2 | 17 | 47 | — | 0 | 2 | 3 | 3 |
| Tennessee† | 1 | 1 | 5 | 60 | 144 | — | 2 | 7 | 122 | 141 | 1 | 1 | 7 | 42 | 36 |
| W.S. Central | 4 | 7 | 77 | 323 | 432 | — | 13 | 315 | 644 | 575 | 6 | 0 | 32 | 49 | 43 |
| Arkansas | — | 0 | 9 | 38 | 18 | — | 1 | 3 | 50 | 65 | — | 0 | 3 | 3 | 6 |
| Louisiana | — | 0 | 4 | 20 | 62 | — | 0 | 5 | 33 | 66 | — | 0 | 2 | 4 | 2 |
| Oklahoma | 3 | 0 | 2 | 9 | 5 | — | 0 | 17 | 70 | 39 | 6 | 0 | 3 | 7 | 7 |
| Texas† | 1 | 5 | 73 | 256 | 347 | — | 10 | 295 | 491 | 405 | — | 0 | 26 | 35 | 28 |
| Mountain | 1 | 5 | 17 | 247 | 302 | 2 | 3 | 16 | 155 | 173 | — | 2 | 8 | 115 | 91 |
| Arizona | — | 2 | 16 | 151 | 169 | — | 0 | 3 | 32 | — | — | 1 | 5 | 38 | 23 |
| Colorado | 1 | 1 | 4 | 36 | 39 | 2 | 1 | 5 | 34 | 53 | — | 0 | 2 | 21 | 19 |
| Idaho† | — | 0 | 2 | 9 | 21 | — | 0 | 2 | 13 | 16 | — | 0 | 3 | 11 | 4 |
| Montana† | — | 0 | 3 | 11 | 9 | — | 0 | 7 | — | 3 | — | 0 | 1 | 6 | 5 |
| Nevada† | — | 0 | 2 | 11 | 20 | — | 1 | 5 | 30 | 46 | — | 0 | 2 | 8 | 19 |
| New Mexico† | — | 0 | 3 | 13 | 24 | — | 0 | 2 | 19 | 18 | — | 0 | 1 | 5 | 4 |
| Utah | — | 0 | 2 | 13 | 19 | — | 0 | 5 | 27 | 35 | — | 0 | 6 | 26 | 13 |
| Wyoming | — | 0 | 1 | 3 | 1 | — | 0 | 1 | — | 2 | — | 0 | 0 | — | 4 |
| Pacific | 7 | 18 | 163 | 982 | 707 | 4 | 11 | 61 | 514 | 511 | 1 | 1 | 9 | 79 | 69 |
| Alaska | — | 0 | 0 | — | 4 | — | 0 | 3 | 9 | 7 | — | 0 | 0 | — | 1 |
| California | 3 | 15 | 162 | 885 | 590 | 1 | 8 | 41 | 381 | 344 | 1 | 1 | 9 | 79 | 65 |
| Hawaii | — | 0 | 2 | 10 | 24 | — | 0 | 1 | 6 | 8 | — | 0 | 0 | — | 3 |
| Oregon† | 1 | 0 | 5 | 40 | 44 | 2 | 1 | 5 | 73 | 93 | N | 0 | 0 | N | N |
| Washington | 3 | 0 | 13 | 47 | 45 | 1 | 0 | 18 | 45 | 59 | — | 0 | 0 | — | — |
| American Samoa | U | 0 | 0 | U | 1 | U | 0 | 0 | U | — | U | 0 | 0 | U | U |
| C.N.M.I. | U | 0 | 0 | U | U | U | 0 | 0 | U | U | U | 0 | 0 | U | U |
| Guam | — | 0 | 0 | — | 2 | — | 0 | 0 | — | 18 | — | 0 | 0 | — | — |
| Puerto Rico | — | 0 | 6 | 30 | 60 | — | 0 | 8 | 27 | 49 | — | 0 | 1 | 1 | — |
| U.S. Virgin Islands | — | 0 | 0 | — | — | — | 0 | 0 | — | — | — | 0 | 0 | — | — |

C.N.M.I.: Commonwealth of Northern Mariana Islands.

U: Unavailable. —: No reported cases. N: Not notifiable.

Cum: Cumulative year-to-date counts.

Med: Median.

Max: Maximum.

* Incidence data for reporting year 2006 is provisional.

† Contains data reported through the National Electronic Disease Surveillance System (NEDSS).

TABLE II. (Continued) Provisional cases of selected notifiable diseases, United States, weeks ending November 25, 2006, and November 26, 2005 (47th Week)*

| Reporting area | Lyme disease | | | | | Malaria | | | | |
|----------------------|--------------|-------------------|-------|----------|----------|--------------|-------------------|-----|----------|----------|
| | Current week | Previous 52 weeks | | Cum 2006 | Cum 2005 | Current week | Previous 52 weeks | | Cum 2006 | Cum 2005 |
| | | Med | Max | | | | Med | Max | | |
| United States | 152 | 236 | 2,153 | 15,780 | 20,175 | 7 | 26 | 125 | 1,145 | 1,269 |
| New England | 83 | 30 | 780 | 2,798 | 3,671 | — | 1 | 11 | 45 | 68 |
| Connecticut | 16 | 11 | 753 | 1,646 | 823 | — | 0 | 3 | 11 | 18 |
| Maine† | 18 | 1 | 34 | 271 | 241 | — | 0 | 1 | 4 | 5 |
| Massachusetts | — | 0 | 14 | 33 | 2,287 | — | 0 | 3 | 19 | 36 |
| New Hampshire | — | 5 | 90 | 522 | 233 | — | 0 | 3 | 9 | 6 |
| Rhode Island | 49 | 0 | 93 | 235 | 37 | — | 0 | 8 | 1 | 2 |
| Vermont† | — | 1 | 14 | 91 | 50 | — | 0 | 1 | 1 | 1 |
| Mid. Atlantic | 57 | 129 | 1,176 | 8,925 | 11,526 | 4 | 5 | 13 | 249 | 333 |
| New Jersey | 2 | 22 | 173 | 1,918 | 3,290 | — | 0 | 3 | 28 | 74 |
| New York (Upstate) | 52 | 59 | 1,150 | 3,734 | 3,728 | 4 | 1 | 11 | 46 | 48 |
| New York City | — | 1 | 18 | 153 | 385 | — | 3 | 9 | 133 | 177 |
| Pennsylvania | 3 | 40 | 235 | 3,120 | 4,123 | — | 1 | 4 | 42 | 34 |
| E.N. Central | — | 9 | 145 | 1,368 | 1,700 | — | 2 | 7 | 113 | 137 |
| Illinois | — | 0 | 1 | — | 126 | — | 1 | 4 | 45 | 71 |
| Indiana | — | 0 | 3 | 19 | 30 | — | 0 | 3 | 10 | 7 |
| Michigan | — | 1 | 6 | 49 | 56 | — | 0 | 2 | 17 | 21 |
| Ohio | — | 1 | 5 | 42 | 53 | — | 0 | 3 | 27 | 24 |
| Wisconsin | — | 8 | 141 | 1,258 | 1,435 | — | 0 | 3 | 14 | 14 |
| W.N. Central | — | 6 | 169 | 720 | 878 | 1 | 0 | 32 | 59 | 46 |
| Iowa | — | 1 | 8 | 87 | 91 | — | 0 | 1 | 2 | 8 |
| Kansas | — | 0 | 2 | 5 | 3 | — | 0 | 2 | 7 | 7 |
| Minnesota | — | 3 | 167 | 606 | 765 | — | 0 | 30 | 37 | 11 |
| Missouri | — | 0 | 2 | 10 | 14 | — | 0 | 1 | 6 | 17 |
| Nebraska† | — | 0 | 2 | 11 | 3 | 1 | 0 | 1 | 5 | 3 |
| North Dakota | — | 0 | 3 | — | — | — | 0 | 1 | 1 | — |
| South Dakota | — | 0 | 1 | 1 | 2 | — | 0 | 1 | 1 | — |
| S. Atlantic | 10 | 26 | 113 | 1,687 | 2,154 | — | 7 | 15 | 294 | 280 |
| Delaware | — | 7 | 28 | 447 | 623 | — | 0 | 1 | 5 | 3 |
| District of Columbia | — | 0 | 7 | 56 | 8 | — | 0 | 2 | 5 | 8 |
| Florida | 4 | 1 | 5 | 49 | 43 | — | 1 | 4 | 56 | 57 |
| Georgia | — | 0 | 1 | 6 | 6 | — | 1 | 6 | 76 | 47 |
| Maryland† | 6 | 13 | 70 | 821 | 1,159 | — | 1 | 5 | 65 | 95 |
| North Carolina | — | 0 | 4 | 29 | 44 | — | 0 | 8 | 28 | 30 |
| South Carolina† | — | 0 | 2 | 18 | 19 | — | 0 | 2 | 9 | 8 |
| Virginia† | — | 3 | 25 | 248 | 235 | — | 1 | 9 | 48 | 29 |
| West Virginia | — | 0 | 44 | 13 | 17 | — | 0 | 1 | 2 | 3 |
| E.S. Central | — | 0 | 3 | 28 | 34 | 1 | 0 | 3 | 22 | 29 |
| Alabama† | — | 0 | 3 | 10 | 3 | — | 0 | 2 | 9 | 6 |
| Kentucky | — | 0 | 2 | 7 | 5 | 1 | 0 | 1 | 4 | 10 |
| Mississippi | — | 0 | 1 | 1 | — | — | 0 | 1 | 4 | — |
| Tennessee† | — | 0 | 2 | 10 | 26 | — | 0 | 2 | 5 | 13 |
| W.S. Central | 1 | 0 | 3 | 18 | 76 | 1 | 2 | 31 | 83 | 116 |
| Arkansas | — | 0 | 1 | — | 4 | — | 0 | 1 | 2 | 6 |
| Louisiana | — | 0 | 0 | — | 3 | — | 0 | 1 | 5 | 5 |
| Oklahoma | — | 0 | 0 | — | — | — | 0 | 2 | 7 | 10 |
| Texas† | 1 | 0 | 3 | 18 | 69 | 1 | 1 | 29 | 69 | 95 |
| Mountain | — | 0 | 4 | 27 | 21 | — | 1 | 9 | 65 | 52 |
| Arizona | — | 0 | 2 | 9 | 8 | — | 0 | 9 | 22 | 13 |
| Colorado | — | 0 | 1 | 1 | — | — | 0 | 2 | 15 | 24 |
| Idaho† | — | 0 | 2 | 6 | 2 | — | 0 | 1 | 1 | — |
| Montana† | — | 0 | 0 | — | — | — | 0 | 1 | 2 | — |
| Nevada† | — | 0 | 1 | 2 | 3 | — | 0 | 1 | 4 | 3 |
| New Mexico† | — | 0 | 1 | 2 | 3 | — | 0 | 1 | 4 | 3 |
| Utah | — | 0 | 1 | 6 | 2 | — | 0 | 2 | 17 | 7 |
| Wyoming | — | 0 | 1 | 1 | 3 | — | 0 | 0 | — | 2 |
| Pacific | 1 | 4 | 16 | 209 | 115 | — | 4 | 13 | 215 | 208 |
| Alaska | — | 0 | 1 | 3 | 4 | — | 0 | 4 | 23 | 6 |
| California | 1 | 4 | 15 | 190 | 81 | — | 4 | 10 | 144 | 153 |
| Hawaii | N | 0 | 0 | N | N | — | 0 | 2 | 4 | 18 |
| Oregon† | — | 0 | 2 | 13 | 20 | — | 0 | 2 | 12 | 13 |
| Washington | — | 0 | 3 | 3 | 10 | — | 0 | 5 | 32 | 18 |
| American Samoa | U | 0 | 0 | U | U | U | 0 | 0 | U | U |
| C.N.M.I. | U | 0 | 0 | U | U | U | 0 | 0 | U | U |
| Guam | — | 0 | 0 | — | — | — | 0 | 0 | — | — |
| Puerto Rico | N | 0 | 0 | N | N | — | 0 | 1 | 1 | 4 |
| U.S. Virgin Islands | — | 0 | 0 | — | — | — | 0 | 0 | — | — |

C.N.M.I.: Commonwealth of Northern Mariana Islands.

U: Unavailable. —: No reported cases. N: Not notifiable.

Cum: Cumulative year-to-date counts.

Med: Median.

Max: Maximum.

*: Incidence data for reporting year 2006 is provisional.

† Contains data reported through the National Electronic Disease Surveillance System (NEDSS).

TABLE II. (Continued) Provisional cases of selected notifiable diseases, United States, weeks ending November 25, 2006, and November 26, 2005 (47th Week)*

| Reporting area | Meningococcal disease, invasive | | | | | | | | | | Pertussis | | | | |
|----------------------|---------------------------------|-------------------|-----|----------|----------|-------------------|-------------------|-----|----------|----------|--------------|-------------------|-------|----------|----------|
| | All serogroups | | | | | Serogroup unknown | | | | | | | | | |
| | Current week | Previous 52 weeks | | Cum 2006 | Cum 2005 | Current week | Previous 52 weeks | | Cum 2006 | Cum 2005 | Current week | Previous 52 weeks | | Cum 2006 | Cum 2005 |
| | | Med | Max | | | | Med | Max | | | | Med | Max | | |
| United States | 10 | 19 | 85 | 907 | 1,091 | 7 | 12 | 58 | 601 | 677 | 67 | 257 | 2,877 | 11,583 | 20,864 |
| New England | 1 | 1 | 3 | 42 | 64 | 1 | 0 | 2 | 29 | 22 | 1 | 25 | 83 | 1,040 | 1,349 |
| Connecticut | — | 0 | 2 | 10 | 12 | — | 0 | 2 | 3 | 1 | — | 1 | 5 | 45 | 67 |
| Maine† | — | 0 | 1 | 6 | 2 | — | 0 | 1 | 4 | 2 | — | 1 | 11 | 84 | 50 |
| Massachusetts | — | 0 | 2 | 15 | 30 | — | 0 | 2 | 15 | 5 | — | 16 | 43 | 594 | 1,016 |
| New Hampshire | — | 0 | 2 | 6 | 12 | — | 0 | 2 | 6 | 12 | — | 2 | 36 | 163 | 97 |
| Rhode Island | — | 0 | 1 | 2 | 3 | — | 0 | 0 | — | — | 1 | 0 | 17 | 50 | 36 |
| Vermont† | 1 | 0 | 1 | 3 | 5 | 1 | 0 | 0 | 1 | 2 | — | 2 | 14 | 104 | 83 |
| Mid. Atlantic | — | 2 | 13 | 100 | 140 | — | 2 | 11 | 96 | 108 | 14 | 36 | 137 | 1,663 | 1,206 |
| New Jersey | N | 0 | 1 | N | 31 | N | 0 | 1 | N | 31 | — | 4 | 13 | 185 | 174 |
| New York (Upstate) | N | 0 | 7 | N | 37 | N | 0 | 5 | N | 13 | 12 | 15 | 123 | 785 | 470 |
| New York City | — | 1 | 4 | 58 | 24 | — | 1 | 4 | 58 | 24 | — | 1 | 8 | 64 | 98 |
| Pennsylvania | — | 1 | 5 | 42 | 48 | — | 1 | 5 | 38 | 40 | 2 | 13 | 26 | 629 | 464 |
| E.N. Central | 3 | 2 | 11 | 108 | 150 | 1 | 1 | 6 | 75 | 118 | 22 | 39 | 133 | 1,725 | 3,547 |
| Illinois | — | 0 | 4 | 18 | 33 | — | 0 | 4 | 18 | 33 | — | 6 | 23 | 231 | 853 |
| Indiana | 1 | 0 | 5 | 22 | 18 | — | 0 | 1 | 8 | 8 | 8 | 4 | 75 | 221 | 298 |
| Michigan | — | 0 | 3 | 20 | 34 | — | 0 | 1 | 9 | 18 | 7 | 9 | 39 | 557 | 291 |
| Ohio | 2 | 1 | 4 | 43 | 42 | 1 | 1 | 3 | 35 | 36 | 7 | 12 | 29 | 548 | 1,062 |
| Wisconsin | — | 0 | 2 | 5 | 23 | — | 0 | 2 | 5 | 23 | — | 4 | 19 | 168 | 1,043 |
| W.N. Central | — | 1 | 4 | 56 | 77 | — | 0 | 3 | 18 | 33 | 3 | 24 | 552 | 1,101 | 3,585 |
| Iowa | — | 0 | 2 | 18 | 15 | — | 0 | 1 | 5 | 1 | — | 6 | 38 | 250 | 1,016 |
| Kansas | — | 0 | 1 | 2 | 9 | — | 0 | 1 | 2 | 9 | 1 | 6 | 25 | 282 | 455 |
| Minnesota | — | 0 | 2 | 13 | 15 | — | 0 | 1 | 4 | 6 | — | 0 | 485 | 161 | 1,025 |
| Missouri | — | 0 | 2 | 14 | 28 | — | 0 | 1 | 2 | 13 | 1 | 6 | 42 | 274 | 500 |
| Nebraska† | — | 0 | 2 | 6 | 5 | — | 0 | 1 | 4 | 3 | 1 | 2 | 9 | 88 | 274 |
| North Dakota | — | 0 | 1 | 1 | 1 | — | 0 | 1 | 1 | 1 | — | 0 | 25 | 26 | 139 |
| South Dakota | — | 0 | 1 | 2 | 4 | — | 0 | 0 | — | — | — | 0 | 4 | 20 | 176 |
| S. Atlantic | 2 | 4 | 14 | 172 | 205 | 1 | 1 | 7 | 72 | 92 | 3 | 18 | 46 | 909 | 1,306 |
| Delaware | — | 0 | 1 | 4 | 4 | — | 0 | 1 | 4 | 4 | — | 0 | 1 | 3 | 15 |
| District of Columbia | 1 | 0 | 1 | 2 | 5 | 1 | 0 | 1 | 2 | 4 | — | 0 | 3 | 6 | 8 |
| Florida | — | 1 | 6 | 65 | 75 | — | 0 | 5 | 24 | 31 | 2 | 4 | 9 | 194 | 187 |
| Georgia | — | 0 | 3 | 14 | 15 | — | 0 | 3 | 14 | 15 | — | 0 | 3 | 22 | 46 |
| Maryland† | — | 0 | 2 | 12 | 22 | — | 0 | 1 | 2 | 5 | 1 | 3 | 9 | 119 | 188 |
| North Carolina | — | 0 | 11 | 30 | 32 | — | 0 | 3 | 10 | 9 | — | 0 | 22 | 177 | 118 |
| South Carolina† | — | 0 | 2 | 20 | 13 | — | 0 | 2 | 9 | 8 | — | 3 | 11 | 162 | 383 |
| Virginia† | — | 0 | 4 | 16 | 33 | — | 0 | 1 | 7 | 14 | — | 1 | 27 | 183 | 316 |
| West Virginia | 1 | 0 | 2 | 9 | 6 | — | 0 | 0 | — | 2 | — | 0 | 9 | 43 | 45 |
| E.S. Central | — | 1 | 4 | 40 | 53 | — | 1 | 4 | 32 | 42 | 3 | 7 | 27 | 347 | 477 |
| Alabama† | — | 0 | 1 | 6 | 5 | — | 0 | 1 | 4 | 3 | 2 | 1 | 18 | 106 | 78 |
| Kentucky | — | 0 | 2 | 11 | 17 | — | 0 | 2 | 11 | 17 | — | 1 | 5 | 54 | 143 |
| Mississippi | — | 0 | 1 | 4 | 7 | — | 0 | 1 | 4 | 7 | 1 | 1 | 4 | 41 | 58 |
| Tennessee† | — | 0 | 2 | 19 | 24 | — | 0 | 2 | 13 | 15 | — | 3 | 10 | 146 | 198 |
| W.S. Central | — | 1 | 23 | 55 | 100 | — | 0 | 6 | 23 | 25 | 7 | 15 | 360 | 673 | 2,179 |
| Arkansas | — | 0 | 3 | 9 | 14 | — | 0 | 2 | 6 | 3 | 2 | 1 | 21 | 75 | 286 |
| Louisiana | — | 0 | 2 | 6 | 29 | — | 0 | 1 | 3 | 6 | — | 0 | 2 | 13 | 49 |
| Oklahoma | — | 0 | 4 | 11 | 14 | — | 0 | 0 | — | 2 | — | 0 | 124 | 19 | 3 |
| Texas† | — | 0 | 16 | 29 | 43 | — | 0 | 4 | 14 | 14 | 5 | 13 | 215 | 566 | 1,841 |
| Mountain | 1 | 1 | 5 | 64 | 82 | 1 | 0 | 4 | 24 | 23 | 8 | 53 | 230 | 2,369 | 3,717 |
| Arizona | — | 0 | 3 | 17 | 31 | — | 0 | 2 | 10 | 10 | — | 8 | 177 | 447 | 896 |
| Colorado | — | 0 | 2 | 20 | 17 | — | 0 | 1 | 2 | — | 6 | 14 | 40 | 703 | 1,234 |
| Idaho† | 1 | 0 | 1 | 4 | 6 | 1 | 0 | 1 | 3 | 5 | 2 | 1 | 8 | 84 | 199 |
| Montana† | — | 0 | 1 | 4 | — | — | 0 | 1 | 2 | — | — | 2 | 9 | 105 | 572 |
| Nevada† | — | 0 | 1 | 4 | 12 | — | 0 | 0 | — | 2 | — | 0 | 9 | 55 | 49 |
| New Mexico† | — | 0 | 1 | 6 | 5 | — | 0 | 1 | 3 | 4 | — | 2 | 6 | 108 | 176 |
| Utah | — | 0 | 1 | 5 | 11 | — | 0 | 0 | — | 2 | — | 15 | 39 | 795 | 542 |
| Wyoming | — | 0 | 2 | 4 | — | — | 0 | 2 | 4 | — | — | 1 | 8 | 72 | 49 |
| Pacific | 3 | 5 | 29 | 270 | 220 | 3 | 5 | 25 | 232 | 214 | 6 | 31 | 1,334 | 1,756 | 3,498 |
| Alaska | 1 | 0 | 1 | 3 | 3 | 1 | 0 | 1 | 3 | 3 | — | 1 | 15 | 63 | 133 |
| California | 1 | 3 | 14 | 167 | 138 | 1 | 3 | 14 | 167 | 138 | — | 22 | 1,136 | 1,249 | 1,769 |
| Hawaii | — | 0 | 1 | 7 | 11 | — | 0 | 1 | 7 | 6 | — | 1 | 4 | 70 | 159 |
| Oregon† | — | 1 | 7 | 62 | 49 | — | 1 | 4 | 43 | 49 | 1 | 1 | 8 | 95 | 616 |
| Washington | 1 | 0 | 25 | 31 | 19 | 1 | 0 | 11 | 12 | 18 | 5 | 4 | 195 | 279 | 821 |
| American Samoa | U | 0 | 0 | — | — | U | 0 | 0 | U | U | U | 0 | 0 | U | U |
| C.N.M.I. | U | 0 | 0 | — | — | U | 0 | 0 | U | U | U | 0 | 0 | U | U |
| Guam | — | 0 | 0 | — | 1 | — | 0 | 0 | — | 1 | — | 0 | 0 | — | 2 |
| Puerto Rico | N | 0 | 0 | — | 7 | N | 0 | 0 | N | 7 | — | 0 | 1 | 2 | 6 |
| U.S. Virgin Islands | — | 0 | 0 | — | — | — | 0 | 0 | — | — | — | 0 | 0 | — | — |

C.N.M.I.: Commonwealth of Northern Mariana Islands.

U: Unavailable. —: No reported cases. N: Not notifiable.

Cum: Cumulative year-to-date counts.

Med: Median.

Max: Maximum.

* Incidence data for reporting year 2006 is provisional.

† Contains data reported through the National Electronic Disease Surveillance System (NEDSS).

TABLE II. (Continued) Provisional cases of selected notifiable diseases, United States, weeks ending November 25, 2006, and November 26, 2005 (47th Week)*

| Reporting area | Rabies, animal | | | | | Rocky Mountain spotted fever | | | | | Salmonellosis | | | | |
|----------------------|----------------|-------------------|-----|----------|----------|------------------------------|-------------------|-----|----------|----------|---------------|-------------------|-------|----------|----------|
| | Current week | Previous 52 weeks | | Cum 2006 | Cum 2005 | Current week | Previous 52 weeks | | Cum 2006 | Cum 2005 | Current week | Previous 52 weeks | | Cum 2006 | Cum 2005 |
| | | Med | Max | | | | Med | Max | | | | Med | Max | | |
| United States | 39 | 119 | 231 | 5,683 | 5,412 | 28 | 38 | 246 | 1,950 | 1,652 | 291 | 779 | 2,291 | 37,606 | 39,972 |
| New England | 5 | 12 | 26 | 615 | 649 | — | 0 | 2 | 2 | 8 | 3 | 23 | 456 | 1,686 | 2,001 |
| Connecticut | 3 | 3 | 14 | 195 | 190 | — | 0 | 0 | — | — | — | 0 | 448 | 448 | 437 |
| Maine† | — | 2 | 8 | 105 | 54 | N | 0 | 0 | N | N | — | 2 | 10 | 109 | 157 |
| Massachusetts | — | 4 | 17 | 178 | 313 | — | 0 | 1 | 1 | 6 | — | 16 | 53 | 782 | 1,059 |
| New Hampshire | 2 | 1 | 5 | 50 | 12 | — | 0 | 1 | 1 | 1 | 1 | 3 | 25 | 195 | 163 |
| Rhode Island | — | 0 | 3 | 24 | 27 | — | 0 | 2 | — | 1 | — | 0 | 17 | 83 | 95 |
| Vermont† | — | 1 | 5 | 63 | 53 | — | 0 | 0 | — | — | 2 | 1 | 6 | 69 | 90 |
| Mid. Atlantic | 7 | 27 | 61 | 1,415 | 916 | 1 | 1 | 5 | 75 | 94 | 27 | 83 | 272 | 4,602 | 4,711 |
| New Jersey | N | 0 | 0 | N | N | — | 0 | 1 | 7 | 29 | — | 14 | 48 | 803 | 912 |
| New York (Upstate) | 5 | 10 | 24 | 504 | 512 | — | 0 | 2 | 5 | 1 | 20 | 24 | 233 | 1,175 | 1,116 |
| New York City | 2 | 0 | 5 | 35 | 28 | — | 0 | 3 | 19 | 7 | 1 | 22 | 50 | 1,116 | 1,123 |
| Pennsylvania | — | 16 | 45 | 876 | 376 | 1 | 1 | 3 | 44 | 57 | 6 | 29 | 67 | 1,508 | 1,560 |
| E.N. Central | 4 | 2 | 18 | 161 | 168 | — | 0 | 6 | 41 | 41 | 36 | 102 | 187 | 4,572 | 5,223 |
| Illinois | — | 0 | 7 | 46 | 50 | — | 0 | 2 | 5 | 11 | — | 23 | 51 | 1,005 | 1,716 |
| Indiana | — | 0 | 2 | 11 | 11 | — | 0 | 1 | 7 | 1 | 6 | 15 | 67 | 785 | 572 |
| Michigan | — | 1 | 5 | 46 | 37 | — | 0 | 1 | 3 | 6 | — | 18 | 35 | 873 | 851 |
| Ohio | 4 | 0 | 9 | 58 | 70 | — | 0 | 4 | 25 | 21 | 30 | 23 | 56 | 1,160 | 1,214 |
| Wisconsin | N | 0 | 0 | N | N | — | 0 | 1 | 1 | 2 | — | 16 | 27 | 749 | 870 |
| W.N. Central | 3 | 6 | 20 | 296 | 305 | 1 | 3 | 15 | 207 | 151 | 15 | 44 | 107 | 2,400 | 2,367 |
| Iowa | — | 1 | 7 | 57 | — | — | 0 | 1 | 5 | 7 | — | 8 | 22 | 403 | 385 |
| Kansas | 2 | 1 | 5 | 76 | 74 | 1 | 0 | 1 | 3 | 5 | 1 | 7 | 16 | 338 | 333 |
| Minnesota | — | 1 | 6 | 39 | 67 | — | 0 | 2 | 4 | 2 | — | 11 | 60 | 650 | 515 |
| Missouri | 1 | 1 | 6 | 65 | 70 | — | 3 | 11 | 171 | 125 | 6 | 14 | 35 | 693 | 740 |
| Nebraska† | — | 0 | 0 | — | — | — | 0 | 5 | 24 | 7 | 7 | 3 | 8 | 175 | 208 |
| North Dakota | — | 0 | 7 | 24 | 30 | — | 0 | 1 | — | — | 1 | 0 | 46 | 28 | 38 |
| South Dakota | — | 1 | 4 | 35 | 64 | — | 0 | 0 | — | 5 | — | 2 | 7 | 113 | 148 |
| S. Atlantic | 17 | 38 | 176 | 2,003 | 1,963 | 25 | 16 | 94 | 1,097 | 831 | 99 | 219 | 392 | 10,208 | 11,660 |
| Delaware | — | 0 | 0 | — | — | — | 0 | 3 | 18 | 7 | — | 2 | 10 | 137 | 116 |
| District of Columbia | — | 0 | 0 | — | — | — | 0 | 1 | 1 | 2 | 2 | 1 | 4 | 59 | 53 |
| Florida | — | 0 | 160 | 160 | 201 | 1 | 0 | 3 | 20 | 13 | 54 | 95 | 176 | 4,318 | 4,832 |
| Georgia | — | 5 | 24 | 213 | 241 | 1 | 0 | 5 | 42 | 85 | 14 | 30 | 72 | 1,586 | 1,819 |
| Maryland† | — | 7 | 13 | 315 | 354 | 1 | 1 | 6 | 71 | 67 | 9 | 12 | 29 | 650 | 759 |
| North Carolina | 11 | 9 | 22 | 481 | 446 | 22 | 14 | 87 | 817 | 468 | 13 | 33 | 130 | 1,521 | 1,556 |
| South Carolina† | — | 3 | 11 | 160 | 206 | — | 0 | 5 | 33 | 71 | — | 18 | 51 | 921 | 1,313 |
| Virginia† | 6 | 11 | 27 | 573 | 450 | — | 1 | 13 | 92 | 111 | 4 | 20 | 57 | 889 | 1,039 |
| West Virginia | — | 2 | 13 | 101 | 65 | — | 0 | 2 | 3 | 7 | 3 | 2 | 19 | 127 | 173 |
| E.S. Central | 2 | 4 | 16 | 226 | 142 | 1 | 5 | 30 | 354 | 284 | 33 | 52 | 149 | 2,827 | 2,751 |
| Alabama† | 1 | 1 | 8 | 79 | 75 | 1 | 1 | 10 | 115 | 72 | 17 | 15 | 71 | 1,005 | 668 |
| Kentucky | 1 | 0 | 4 | 28 | 17 | — | 0 | 1 | 3 | 3 | 3 | 8 | 23 | 406 | 456 |
| Mississippi | — | 0 | 2 | 4 | 5 | — | 0 | 1 | 4 | 18 | — | 11 | 42 | 709 | 863 |
| Tennessee† | — | 2 | 9 | 115 | 45 | — | 3 | 21 | 232 | 191 | 13 | 14 | 31 | 707 | 764 |
| W.S. Central | — | 11 | 34 | 562 | 812 | — | 1 | 161 | 115 | 209 | 15 | 74 | 922 | 3,724 | 3,984 |
| Arkansas | — | 0 | 5 | 31 | 33 | — | 0 | 10 | 51 | 121 | 7 | 15 | 47 | 865 | 686 |
| Louisiana | — | 0 | 0 | — | — | — | 0 | 1 | 4 | 6 | — | 12 | 42 | 740 | 859 |
| Oklahoma | — | 1 | 9 | 60 | 74 | — | 0 | 154 | 36 | 52 | 8 | 8 | 48 | 462 | 377 |
| Texas† | — | 10 | 29 | 471 | 705 | — | 0 | 4 | 24 | 30 | — | 31 | 839 | 1,657 | 2,062 |
| Mountain | — | 3 | 27 | 199 | 254 | — | 0 | 6 | 52 | 32 | 15 | 52 | 88 | 2,322 | 2,201 |
| Arizona | — | 2 | 10 | 129 | 165 | — | 0 | 6 | 13 | 17 | 5 | 17 | 67 | 786 | 616 |
| Colorado | — | 0 | 0 | — | 18 | — | 0 | 1 | 2 | 4 | 8 | 12 | 30 | 565 | 536 |
| Idaho† | — | 0 | 25 | 25 | — | — | 0 | 3 | 14 | 3 | 2 | 3 | 9 | 161 | 141 |
| Montana† | — | 0 | 2 | 14 | 15 | — | 0 | 2 | 2 | 1 | — | 3 | 16 | 118 | 125 |
| Nevada† | — | 0 | 1 | 2 | 14 | — | 0 | 0 | — | — | — | 3 | 20 | 174 | 183 |
| New Mexico† | — | 0 | 2 | 10 | 10 | — | 0 | 2 | 8 | 4 | — | 4 | 15 | 221 | 232 |
| Utah | — | 0 | 1 | 11 | 15 | — | 0 | 2 | 6 | — | — | 5 | 15 | 254 | 287 |
| Wyoming | — | 0 | 2 | 8 | 17 | — | 0 | 1 | 7 | 3 | — | 1 | 4 | 43 | 81 |
| Pacific | 1 | 4 | 12 | 206 | 203 | — | 0 | 1 | 7 | 2 | 48 | 111 | 426 | 5,265 | 5,074 |
| Alaska | — | 0 | 4 | 15 | 1 | — | 0 | 0 | — | — | 1 | 1 | 7 | 67 | 57 |
| California | 1 | 3 | 11 | 166 | 195 | — | 0 | 1 | 5 | — | 41 | 90 | 292 | 4,161 | 3,867 |
| Hawaii | — | 0 | 0 | — | — | — | 0 | 0 | — | — | — | 5 | 10 | 220 | 273 |
| Oregon† | — | 0 | 4 | 25 | 7 | — | 0 | 1 | 2 | 2 | 1 | 8 | 16 | 373 | 385 |
| Washington | U | 0 | 0 | U | U | N | 0 | 0 | N | N | 5 | 8 | 124 | 444 | 492 |
| American Samoa | U | 0 | 0 | U | U | U | 0 | 0 | U | U | U | 0 | 0 | U | 7 |
| C.N.M.I. | U | 0 | 0 | U | U | U | 0 | 0 | U | U | U | 0 | 0 | U | U |
| Guam | — | 0 | 0 | — | — | — | 0 | 0 | — | — | — | 2 | 3 | — | 37 |
| Puerto Rico | — | 1 | 6 | 68 | 61 | N | 0 | 0 | N | N | — | 4 | 35 | 230 | 586 |
| U.S. Virgin Islands | — | 0 | 0 | — | — | — | 0 | 0 | — | — | — | 0 | 0 | — | — |

C.N.M.I.: Commonwealth of Northern Mariana Islands.

U: Unavailable. —: No reported cases. N: Not notifiable.

Cum: Cumulative year-to-date counts.

Med: Median.

Max: Maximum.

* Incidence data for reporting year 2006 is provisional.

† Contains data reported through the National Electronic Disease Surveillance System (NEDSS).

TABLE II. (Continued) Provisional cases of selected notifiable diseases, United States, weeks ending November 25, 2006, and November 26, 2005 (47th Week)*

| Reporting area | Shiga toxin-producing <i>E. coli</i> (STEC) [†] | | | | | Shigellosis | | | | | Streptococcal disease, invasive, group A | | | | |
|-----------------------------|--|-------------------|-----|----------|----------|--------------|-------------------|-------|----------|----------|--|-------------------|-----|----------|----------|
| | Current week | Previous 52 weeks | | Cum 2006 | Cum 2005 | Current week | Previous 52 weeks | | Cum 2006 | Cum 2005 | Current week | Previous 52 weeks | | Cum 2006 | Cum 2005 |
| | | Med | Max | | | | Med | Max | | | | Med | Max | | |
| United States | 37 | 53 | 297 | 2,548 | 2,987 | 116 | 256 | 1,013 | 12,036 | 13,825 | 26 | 92 | 282 | 4,281 | 4,060 |
| New England | 5 | 3 | 73 | 247 | 208 | — | 3 | 65 | 217 | 297 | 1 | 4 | 15 | 184 | 262 |
| Connecticut | — | 0 | 72 | 72 | 55 | — | 0 | 59 | 59 | 52 | U | 0 | 2 | U | 93 |
| Maine [§] | 5 | 0 | 8 | 43 | 29 | — | 0 | 2 | 3 | 14 | — | 0 | 2 | 17 | 14 |
| Massachusetts | — | 1 | 9 | 82 | 83 | — | 2 | 11 | 128 | 181 | — | 2 | 6 | 101 | 119 |
| New Hampshire | — | 0 | 3 | 25 | 16 | — | 0 | 4 | 8 | 13 | — | 0 | 9 | 44 | 17 |
| Rhode Island | — | 0 | 2 | 8 | 7 | — | 0 | 3 | 13 | 20 | 1 | 0 | 3 | 8 | 9 |
| Vermont [§] | — | 0 | 2 | 2 | 18 | — | 0 | 2 | 6 | 17 | — | 0 | 2 | 14 | 10 |
| Mid. Atlantic | 1 | 4 | 107 | 188 | 334 | 4 | 16 | 72 | 756 | 1,147 | 5 | 18 | 43 | 821 | 799 |
| New Jersey | — | 0 | 3 | 3 | 71 | — | 3 | 34 | 242 | 293 | — | 2 | 8 | 122 | 169 |
| New York (Upstate) | — | 0 | 103 | 10 | 126 | 3 | 4 | 60 | 209 | 242 | 3 | 5 | 32 | 278 | 221 |
| New York City | — | 0 | 4 | 32 | 17 | — | 5 | 13 | 222 | 382 | — | 3 | 8 | 136 | 158 |
| Pennsylvania | — | 0 | 4 | 8 | 120 | 1 | 1 | 6 | 83 | 230 | 2 | 6 | 13 | 285 | 251 |
| E.N. Central | 7 | 10 | 56 | 593 | 604 | 4 | 20 | 37 | 915 | 1,081 | 4 | 14 | 44 | 715 | 820 |
| Illinois | — | 1 | 7 | 75 | 135 | — | 7 | 18 | 316 | 371 | — | 3 | 11 | 144 | 273 |
| Indiana | 1 | 1 | 8 | 78 | 68 | 2 | 2 | 18 | 150 | 167 | — | 2 | 11 | 104 | 93 |
| Michigan | — | 1 | 7 | 86 | 86 | — | 3 | 8 | 139 | 218 | — | 3 | 12 | 196 | 192 |
| Ohio | 6 | 3 | 18 | 179 | 163 | 2 | 3 | 14 | 176 | 111 | 4 | 4 | 19 | 219 | 177 |
| Wisconsin | — | 2 | 39 | 175 | 152 | — | 3 | 9 | 134 | 214 | — | 1 | 4 | 52 | 85 |
| W.N. Central | 4 | 9 | 32 | 495 | 497 | 1 | 35 | 77 | 1,534 | 1,564 | — | 5 | 57 | 313 | 255 |
| Iowa | — | 2 | 8 | 116 | 94 | 1 | 2 | 10 | 103 | 93 | N | 0 | 0 | N | N |
| Kansas | — | 0 | 4 | 25 | 53 | — | 3 | 20 | 133 | 228 | — | 1 | 5 | 52 | 38 |
| Minnesota | — | 3 | 27 | 219 | 163 | — | 3 | 23 | 203 | 83 | — | 0 | 52 | 143 | 96 |
| Missouri | — | 1 | 10 | 82 | 90 | — | 10 | 69 | 613 | 928 | — | 1 | 5 | 71 | 64 |
| Nebraska [§] | — | 1 | 8 | 55 | 58 | — | 2 | 14 | 119 | 137 | — | 0 | 4 | 28 | 22 |
| North Dakota | — | 0 | 15 | — | 8 | — | 0 | 18 | 103 | 4 | — | 0 | 5 | 11 | 10 |
| South Dakota | — | 0 | 5 | 47 | 31 | — | 5 | 22 | 260 | 91 | — | 0 | 1 | 8 | 25 |
| S. Atlantic | 6 | 9 | 39 | 434 | 380 | 65 | 57 | 137 | 2,978 | 2,196 | 6 | 21 | 44 | 1,043 | 849 |
| Delaware | — | 0 | 2 | 9 | 9 | — | 0 | 2 | 10 | 11 | — | 0 | 2 | 10 | 6 |
| District of Columbia | 1 | 0 | 1 | 3 | 1 | 1 | 0 | 2 | 16 | 13 | — | 0 | 2 | 15 | 10 |
| Florida | 3 | 2 | 29 | 87 | 85 | 25 | 27 | 76 | 1,423 | 1,070 | 1 | 5 | 16 | 273 | 229 |
| Georgia | 1 | 2 | 6 | 83 | 49 | 25 | 19 | 73 | 1,092 | 609 | 4 | 5 | 12 | 220 | 186 |
| Maryland [§] | 1 | 1 | 8 | 91 | 72 | 4 | 2 | 10 | 120 | 95 | — | 4 | 12 | 182 | 162 |
| North Carolina | — | 2 | 7 | 104 | 60 | 8 | 1 | 21 | 151 | 184 | — | 0 | 26 | 148 | 118 |
| South Carolina [§] | — | 0 | 2 | 9 | 11 | — | 1 | 9 | 72 | 96 | — | 1 | 6 | 54 | 33 |
| Virginia [§] | — | 0 | 8 | — | 89 | 2 | 1 | 9 | 90 | 117 | 1 | 2 | 11 | 115 | 83 |
| West Virginia | — | 0 | 5 | 12 | 4 | — | 0 | 2 | 4 | 1 | — | 0 | 6 | 26 | 22 |
| E.S. Central | — | 1 | 12 | 92 | 172 | 1 | 13 | 79 | 812 | 1,126 | 1 | 3 | 11 | 179 | 164 |
| Alabama [§] | — | 0 | 5 | 39 | 29 | — | 4 | 71 | 354 | 211 | N | 0 | 0 | N | N |
| Kentucky | — | 1 | 12 | 92 | 74 | 1 | 4 | 15 | 226 | 300 | — | 0 | 5 | 35 | 31 |
| Mississippi | — | 0 | 0 | — | 8 | — | 1 | 9 | 86 | 91 | — | 0 | 0 | — | — |
| Tennessee [§] | — | 0 | 4 | 24 | 61 | — | 3 | 12 | 146 | 524 | 1 | 3 | 9 | 144 | 133 |
| W.S. Central | 8 | 1 | 52 | 76 | 103 | 12 | 36 | 596 | 1,640 | 3,302 | 1 | 7 | 58 | 335 | 285 |
| Arkansas | — | 0 | 7 | 33 | 13 | 3 | 2 | 9 | 113 | 57 | — | 0 | 5 | 25 | 21 |
| Louisiana | — | 0 | 1 | — | 21 | — | 1 | 25 | 132 | 133 | — | 0 | 2 | 8 | — |
| Oklahoma | 8 | 0 | 17 | 43 | 26 | 3 | 2 | 286 | 125 | 602 | — | 2 | 14 | 93 | 105 |
| Texas [§] | 1 | 2 | 44 | 105 | 43 | 6 | 30 | 308 | 1,270 | 2,510 | 1 | 4 | 43 | 209 | 159 |
| Mountain | 3 | 5 | 16 | 297 | 295 | 16 | 23 | 88 | 1,314 | 872 | 6 | 11 | 77 | 578 | 524 |
| Arizona | 3 | 2 | 13 | 119 | 30 | 8 | 13 | 36 | 665 | 459 | 4 | 6 | 57 | 314 | 224 |
| Colorado | — | 1 | 8 | 101 | 79 | 7 | 3 | 15 | 225 | 156 | 2 | 3 | 8 | 123 | 160 |
| Idaho [§] | 3 | 1 | 7 | 79 | 49 | 1 | 0 | 3 | 15 | 17 | — | 0 | 2 | 8 | 3 |
| Montana [§] | — | 0 | 1 | — | 16 | — | 0 | 10 | 41 | 5 | — | 0 | 0 | — | — |
| Nevada [§] | — | 0 | 5 | 22 | 23 | — | 1 | 20 | 103 | 59 | — | 0 | 0 | — | — |
| New Mexico [§] | — | 0 | 1 | 4 | 24 | — | 2 | 15 | 158 | 129 | — | 1 | 7 | 66 | 76 |
| Utah | — | 1 | 14 | 114 | 64 | — | 1 | 6 | 75 | 42 | — | 1 | 7 | 63 | 56 |
| Wyoming | — | 0 | 3 | 18 | 10 | — | 0 | 8 | 32 | 5 | — | 0 | 1 | 4 | 5 |
| Pacific | 3 | 2 | 50 | 126 | 394 | 13 | 39 | 148 | 1,870 | 2,240 | 2 | 2 | 9 | 113 | 102 |
| Alaska | — | 0 | 0 | — | — | — | 0 | 2 | 9 | 11 | — | 0 | 0 | — | — |
| California | — | 2 | 18 | — | 135 | 11 | 32 | 104 | 1,573 | 1,942 | — | 0 | 0 | — | — |
| Hawaii | — | 0 | 2 | 17 | 13 | — | 1 | 4 | 42 | 32 | — | 2 | 9 | 113 | 102 |
| Oregon [§] | — | 2 | 14 | 106 | 152 | — | 1 | 31 | 112 | 121 | N | 0 | 0 | N | N |
| Washington | 3 | 2 | 32 | 109 | 94 | 2 | 2 | 43 | 134 | 134 | N | 0 | 0 | N | N |
| American Samoa | U | 0 | 0 | U | U | U | 0 | 0 | U | 7 | U | 0 | 0 | U | U |
| C.N.M.I. | U | 0 | 0 | U | U | U | 0 | 0 | U | U | U | 0 | 0 | U | U |
| Guam | — | 0 | 0 | — | — | — | 0 | 3 | — | 17 | — | 0 | 0 | — | — |
| Puerto Rico | — | 0 | 0 | — | 2 | — | 0 | 2 | 13 | 9 | N | 0 | 0 | N | N |
| U.S. Virgin Islands | — | 0 | 0 | — | — | — | 0 | 0 | — | — | — | 0 | 0 | — | — |

C.N.M.I.: Commonwealth of Northern Mariana Islands.

U: Unavailable. —: No reported cases. N: Not notifiable.

Cum: Cumulative year-to-date counts.

Med: Median.

Max: Maximum.

* Incidence data for reporting year 2006 is provisional.

† Includes *E. coli* O157:H7; Shiga toxin positive, serogroup non-O157; and Shiga toxin positive, not serogrouped.

§ Contains data reported through the National Electronic Disease Surveillance System (NEDSS).

TABLE II. (Continued) Provisional cases of selected notifiable diseases, United States, weeks ending November 25, 2006, and November 26, 2005 (47th Week)*

| Reporting area | <i>Streptococcus pneumoniae</i> , invasive disease Drug resistant, all ages | | | | | Syphilis, primary and secondary | | | | | Varicella (chickenpox) | | | | |
|----------------------|--|-------------------|-----|----------|----------|---------------------------------|-------------------|-----|----------|----------|------------------------|-------------------|-------|----------|----------|
| | Current week | Previous 52 weeks | | Cum 2006 | Cum 2005 | Current week | Previous 52 weeks | | Cum 2006 | Cum 2005 | Current week | Previous 52 weeks | | Cum 2006 | Cum 2005 |
| | | Med | Max | | | | Med | Max | | | | Med | Max | | |
| United States | 29 | 51 | 333 | 2,222 | 2,293 | 44 | 175 | 334 | 8,118 | 7,685 | 570 | 835 | 2,857 | 37,226 | 26,018 |
| New England | 3 | 1 | 24 | 36 | 206 | 12 | 4 | 17 | 187 | 193 | 27 | 34 | 144 | 1,330 | 4,789 |
| Connecticut | U | 0 | 7 | U | 82 | 10 | 0 | 11 | 48 | 44 | U | 0 | 55 | U | 1,490 |
| Maine† | — | 0 | 2 | 9 | N | — | 0 | 2 | 8 | 1 | — | 2 | 20 | 151 | 277 |
| Massachusetts | — | 0 | 6 | — | 95 | — | 2 | 6 | 107 | 111 | — | 0 | 54 | 94 | 2,094 |
| New Hampshire | — | 0 | 0 | — | — | — | 0 | 2 | 11 | 15 | 4 | 6 | 47 | 447 | 303 |
| Rhode Island | 3 | 0 | 11 | 13 | 18 | 2 | 0 | 2 | 11 | 21 | — | 0 | 0 | — | — |
| Vermont† | — | 0 | 2 | 14 | 11 | — | 0 | 1 | 2 | 1 | 23 | 12 | 50 | 638 | 625 |
| Mid. Atlantic | 4 | 3 | 15 | 158 | 189 | 8 | 21 | 35 | 1,004 | 927 | 86 | 102 | 183 | 4,414 | 4,346 |
| New Jersey | N | 0 | 0 | N | N | 4 | 3 | 8 | 150 | 120 | — | 0 | 0 | — | — |
| New York (Upstate) | 4 | 1 | 10 | 60 | 72 | 2 | 3 | 14 | 136 | 69 | — | 0 | 0 | — | — |
| New York City | U | 0 | 0 | U | U | 1 | 10 | 23 | 488 | 557 | — | 0 | 0 | — | — |
| Pennsylvania | — | 2 | 9 | 98 | 117 | 1 | 5 | 12 | 230 | 181 | 86 | 102 | 183 | 4,414 | 4,346 |
| E.N. Central | 4 | 11 | 41 | 512 | 573 | 4 | 17 | 39 | 805 | 834 | 284 | 245 | 587 | 13,279 | 5,323 |
| Illinois | — | 0 | 3 | 17 | 32 | 1 | 8 | 23 | 381 | 470 | — | 1 | 7 | 68 | 91 |
| Indiana | — | 2 | 21 | 146 | 173 | — | 1 | 4 | 80 | 57 | — | 0 | 475 | 475 | — |
| Michigan | — | 0 | 4 | 18 | 40 | — | 2 | 19 | 109 | 75 | 46 | 102 | 168 | 4,137 | 3,410 |
| Ohio | 4 | 6 | 32 | 331 | 328 | 3 | 4 | 8 | 174 | 197 | 238 | 129 | 420 | 7,955 | 1,414 |
| Wisconsin | N | 0 | 0 | N | N | — | 1 | 4 | 61 | 35 | — | 13 | 52 | 644 | 408 |
| W.N. Central | — | 1 | 191 | 101 | 42 | 1 | 5 | 11 | 236 | 235 | 28 | 28 | 98 | 1,610 | 570 |
| Iowa | N | 0 | 0 | N | N | — | 0 | 3 | 18 | 8 | N | 0 | 0 | N | N |
| Kansas | N | 0 | 0 | N | N | — | 0 | 3 | 23 | 17 | 2 | 3 | 24 | 295 | — |
| Minnesota | — | 0 | 191 | 60 | — | — | 0 | 2 | 26 | 67 | — | 0 | 0 | — | — |
| Missouri | — | 1 | 3 | 39 | 34 | 1 | 3 | 8 | 153 | 137 | 26 | 22 | 82 | 1,196 | 380 |
| Nebraska† | — | 0 | 1 | 1 | 2 | — | 0 | 1 | 3 | 4 | — | 0 | 0 | — | — |
| North Dakota | — | 0 | 1 | — | 3 | — | 0 | 1 | 1 | 1 | — | 0 | 25 | 45 | 61 |
| South Dakota | — | 0 | 1 | 1 | 3 | — | 0 | 3 | 12 | 1 | — | 1 | 10 | 74 | 129 |
| S. Atlantic | 17 | 25 | 53 | 1,173 | 963 | 11 | 42 | 186 | 1,924 | 1,922 | 44 | 88 | 860 | 3,943 | 2,278 |
| Delaware | — | 0 | 0 | — | 3 | — | 0 | 2 | 17 | 10 | — | 1 | 6 | 62 | 28 |
| District of Columbia | — | 0 | 3 | 26 | 13 | — | 2 | 9 | 116 | 102 | — | 0 | 5 | 45 | 37 |
| Florida | 13 | 13 | 36 | 649 | 514 | 5 | 15 | 23 | 670 | 644 | — | 0 | 0 | — | — |
| Georgia | 2 | 7 | 29 | 395 | 325 | — | 7 | 147 | 347 | 442 | — | 0 | 0 | — | — |
| Maryland† | — | 0 | 0 | — | — | — | 5 | 19 | 262 | 273 | — | 0 | 4 | 11 | — |
| North Carolina | N | 0 | 0 | N | N | 1 | 5 | 17 | 272 | 248 | — | 0 | 0 | — | — |
| South Carolina† | — | 0 | 0 | — | — | — | 1 | 6 | 61 | 75 | 6 | 16 | 53 | 962 | 557 |
| Virginia† | N | 0 | 0 | N | N | 5 | 3 | 17 | 174 | 125 | 20 | 28 | 812 | 1,505 | 632 |
| West Virginia | 2 | 1 | 14 | 103 | 108 | — | 0 | 1 | 5 | 3 | 18 | 25 | 70 | 1,358 | 1,024 |
| E.S. Central | 1 | 3 | 13 | 133 | 163 | 1 | 13 | 26 | 663 | 429 | — | 1 | 70 | 119 | 221 |
| Alabama† | N | 0 | 0 | N | N | 1 | 5 | 19 | 288 | 142 | — | 1 | 70 | 117 | 221 |
| Kentucky | — | 0 | 2 | — | 29 | — | 1 | 8 | 63 | 47 | N | 0 | 0 | N | N |
| Mississippi | — | 0 | 0 | — | 1 | — | 1 | 7 | 69 | 43 | — | 0 | 1 | 2 | — |
| Tennessee† | 1 | 3 | 13 | 133 | 133 | — | 5 | 13 | 243 | 197 | N | 0 | 0 | N | N |
| W.S. Central | — | 0 | 5 | 20 | 107 | 1 | 28 | 52 | 1,412 | 1,140 | 81 | 187 | 1,757 | 10,021 | 6,115 |
| Arkansas | — | 0 | 3 | 12 | 13 | — | 1 | 6 | 74 | 46 | 60 | 9 | 110 | 805 | 25 |
| Louisiana | — | 0 | 4 | 8 | 94 | 1 | 4 | 27 | 264 | 256 | — | 0 | 8 | 48 | 120 |
| Oklahoma | N | 0 | 0 | N | N | — | 1 | 6 | 66 | 36 | — | 0 | 0 | — | — |
| Texas† | N | 0 | 0 | N | N | — | 22 | 36 | 1,008 | 802 | 21 | 170 | 1,647 | 9,168 | 5,970 |
| Mountain | — | 2 | 9 | 89 | 50 | 3 | 8 | 25 | 374 | 384 | 20 | 58 | 137 | 2,510 | 2,376 |
| Arizona | N | 0 | 0 | N | N | 3 | 3 | 16 | 164 | 157 | — | 0 | 0 | — | — |
| Colorado | N | 0 | 0 | N | N | — | 1 | 3 | 44 | 43 | 19 | 31 | 76 | 1,358 | 1,655 |
| Idaho† | N | 0 | 0 | N | N | — | 0 | 1 | 2 | 20 | — | 0 | 0 | — | — |
| Montana† | — | 0 | 1 | — | — | — | 0 | 1 | 1 | 6 | — | 0 | 2 | 2 | — |
| Nevada† | — | 0 | 0 | — | — | — | 1 | 12 | 95 | 98 | — | 0 | 0 | — | — |
| New Mexico† | — | 0 | 1 | 1 | — | — | 1 | 5 | 59 | 51 | 1 | 4 | 34 | 339 | 203 |
| Utah | — | 1 | 9 | 46 | 25 | — | 0 | 2 | 9 | 9 | — | 13 | 55 | 758 | 465 |
| Wyoming | — | 1 | 4 | 42 | 25 | — | 0 | 0 | — | — | — | 0 | 11 | 53 | 53 |
| Pacific | — | 0 | 0 | — | — | 3 | 34 | 51 | 1,513 | 1,621 | — | 0 | 0 | — | — |
| Alaska | — | 0 | 0 | — | — | — | 0 | 4 | 9 | 6 | — | 0 | 0 | — | — |
| California | N | 0 | 0 | N | N | 1 | 29 | 42 | 1,310 | 1,435 | — | 0 | 0 | — | — |
| Hawaii | — | 0 | 0 | — | — | — | 0 | 2 | 17 | 10 | N | 0 | 0 | N | N |
| Oregon† | N | 0 | 0 | N | N | 1 | 0 | 3 | 18 | 33 | N | 0 | 0 | N | N |
| Washington | N | 0 | 0 | N | N | 1 | 2 | 10 | 159 | 137 | N | 0 | 0 | N | N |
| American Samoa | — | 0 | 0 | — | — | U | 0 | 0 | U | U | U | 0 | 0 | U | U |
| C.N.M.I. | — | 0 | 0 | — | — | U | 0 | 0 | U | U | U | 0 | 0 | U | U |
| Guam | — | 0 | 0 | — | — | — | 0 | 0 | — | 3 | — | 2 | 5 | — | 430 |
| Puerto Rico | N | 0 | 0 | N | N | — | 3 | 10 | 120 | 199 | — | 7 | 47 | 316 | 644 |
| U.S. Virgin Islands | — | 0 | 0 | — | — | — | 0 | 0 | — | — | — | 0 | 0 | — | — |

C.N.M.I.: Commonwealth of Northern Mariana Islands.

U: Unavailable. —: No reported cases. N: Not notifiable.

Cum: Cumulative year-to-date counts.

Med: Median.

Max: Maximum.

* Incidence data for reporting year 2006 is provisional.

† Contains data reported through the National Electronic Disease Surveillance System (NEDSS).

TABLE II. (Continued) Provisional cases of selected notifiable diseases, United States, weeks ending November 25, 2006, and November 26, 2005 (47th Week)*

| Reporting area | West Nile virus disease [†] | | | | | | | | | |
|-----------------------------|--------------------------------------|-------------------|-----|----------|----------|-------------------|-------------------|-----|----------|----------|
| | Neuroinvasive | | | | | Non-neuroinvasive | | | | |
| | Current week | Previous 52 weeks | | Cum 2006 | Cum 2005 | Current week | Previous 52 weeks | | Cum 2006 | Cum 2005 |
| | | Med | Max | | | | Med | Max | | |
| United States | — | 1 | 176 | 1,381 | 1,191 | — | 1 | 383 | 2,443 | 1,683 |
| New England | — | 0 | 3 | 9 | 9 | — | 0 | 2 | 3 | 4 |
| Connecticut | — | 0 | 3 | 7 | 4 | — | 0 | 1 | 2 | 2 |
| Maine [§] | — | 0 | 0 | — | — | — | 0 | 0 | — | — |
| Massachusetts | — | 0 | 1 | 2 | 4 | — | 0 | 1 | 1 | 2 |
| New Hampshire | — | 0 | 0 | — | — | — | 0 | 0 | — | — |
| Rhode Island | — | 0 | 0 | — | 1 | — | 0 | 0 | — | — |
| Vermont [§] | — | 0 | 0 | — | — | — | 0 | 0 | — | — |
| Mid. Atlantic | — | 0 | 11 | 26 | 47 | — | 0 | 4 | 10 | 22 |
| New Jersey | — | 0 | 2 | 2 | 3 | — | 0 | 1 | 2 | 3 |
| New York (Upstate) | — | 0 | 5 | 8 | 19 | — | 0 | 1 | 3 | 5 |
| New York City | — | 0 | 4 | 8 | 11 | — | 0 | 2 | 4 | 3 |
| Pennsylvania | — | 0 | 2 | 8 | 14 | — | 0 | 1 | 1 | 11 |
| E.N. Central | — | 0 | 43 | 235 | 259 | — | 0 | 22 | 99 | 156 |
| Illinois | — | 0 | 21 | 116 | 137 | — | 0 | 19 | 70 | 115 |
| Indiana | — | 0 | 7 | 26 | 11 | — | 0 | 2 | 7 | 12 |
| Michigan | — | 0 | 10 | 46 | 54 | — | 0 | 1 | 2 | 8 |
| Ohio | — | 0 | 11 | 36 | 46 | — | 0 | 3 | 11 | 15 |
| Wisconsin | — | 0 | 2 | 11 | 11 | — | 0 | 2 | 9 | 6 |
| W.N. Central | — | 0 | 35 | 216 | 169 | — | 0 | 79 | 473 | 463 |
| Iowa | — | 0 | 3 | 21 | 14 | — | 0 | 4 | 13 | 23 |
| Kansas | — | 0 | 3 | 17 | 17 | — | 0 | 3 | 13 | N |
| Minnesota | — | 0 | 6 | 30 | 18 | — | 0 | 7 | 35 | 27 |
| Missouri | — | 0 | 13 | 47 | 17 | — | 0 | 2 | 12 | 13 |
| Nebraska [§] | — | 0 | 9 | 43 | 55 | — | 0 | 37 | 208 | 133 |
| North Dakota | — | 0 | 5 | 20 | 12 | — | 0 | 28 | 117 | 74 |
| South Dakota | — | 0 | 7 | 38 | 36 | — | 0 | 22 | 75 | 193 |
| S. Atlantic | — | 0 | 2 | 14 | 34 | — | 0 | 4 | 7 | 29 |
| Delaware | — | 0 | 0 | — | 1 | — | 0 | 0 | — | 1 |
| District of Columbia | — | 0 | 0 | — | 3 | — | 0 | 1 | 1 | 2 |
| Florida | — | 0 | 1 | 3 | 10 | — | 0 | 0 | — | 11 |
| Georgia | — | 0 | 1 | 2 | 9 | — | 0 | 3 | 5 | 11 |
| Maryland [§] | — | 0 | 2 | 7 | 4 | — | 0 | 1 | 1 | 1 |
| North Carolina | — | 0 | 0 | — | 2 | — | 0 | 0 | — | 2 |
| South Carolina [§] | — | 0 | 1 | 1 | 5 | — | 0 | 0 | — | — |
| Virginia [§] | — | 0 | 0 | — | — | — | 0 | 0 | — | 1 |
| West Virginia | — | 0 | 1 | 1 | — | N | 0 | 0 | N | N |
| E.S. Central | — | 0 | 14 | 106 | 65 | — | 0 | 15 | 92 | 38 |
| Alabama [§] | — | 0 | 2 | 7 | 6 | — | 0 | 0 | — | 4 |
| Kentucky | — | 0 | 0 | — | 5 | — | 0 | 1 | 1 | — |
| Mississippi | — | 0 | 10 | 84 | 39 | — | 0 | 15 | 89 | 31 |
| Tennessee [§] | — | 0 | 4 | 15 | 15 | — | 0 | 2 | 2 | 3 |
| W.S. Central | — | 0 | 59 | 347 | 157 | — | 0 | 26 | 207 | 150 |
| Arkansas | — | 0 | 4 | 23 | 13 | — | 0 | 2 | 5 | 15 |
| Louisiana | — | 0 | 14 | 88 | — | — | 0 | 9 | 81 | 54 |
| Oklahoma | — | 0 | 6 | 26 | 17 | — | 0 | 4 | 18 | 14 |
| Texas [§] | — | 0 | 38 | 210 | 127 | — | 0 | 15 | 103 | 67 |
| Mountain | — | 0 | 61 | 342 | 145 | — | 0 | 222 | 1,320 | 240 |
| Arizona | — | 0 | 9 | 48 | 52 | — | 0 | 12 | 57 | 61 |
| Colorado | — | 0 | 10 | 63 | 21 | — | 0 | 51 | 269 | 85 |
| Idaho [§] | — | 0 | 30 | 111 | 3 | — | 0 | 151 | 752 | 10 |
| Montana [§] | — | 0 | 3 | 12 | 8 | — | 0 | 7 | 21 | 17 |
| Nevada [§] | — | 0 | 9 | 34 | 14 | — | 0 | 13 | 75 | 17 |
| New Mexico [§] | — | 0 | 1 | 3 | 20 | — | 0 | 1 | 5 | 13 |
| Utah | — | 0 | 8 | 56 | 21 | — | 0 | 17 | 101 | 31 |
| Wyoming | — | 0 | 7 | 15 | 6 | — | 0 | 8 | 40 | 6 |
| Pacific | — | 0 | 15 | 86 | 306 | — | 0 | 45 | 232 | 581 |
| Alaska | — | 0 | 0 | — | — | — | 0 | 0 | — | — |
| California | — | 0 | 15 | 79 | 305 | — | 0 | 33 | 179 | 575 |
| Hawaii | — | 0 | 0 | — | — | — | 0 | 0 | — | — |
| Oregon [§] | — | 0 | 2 | 7 | 1 | — | 0 | 12 | 50 | 6 |
| Washington | — | 0 | 0 | — | — | — | 0 | 2 | 3 | — |
| American Samoa | U | 0 | 0 | U | U | U | 0 | 0 | U | U |
| C.N.M.I. | U | 0 | 0 | U | U | U | 0 | 0 | U | U |
| Guam | — | 0 | 0 | — | — | — | 0 | 0 | — | — |
| Puerto Rico | — | 0 | 0 | — | — | — | 0 | 0 | — | — |
| U.S. Virgin Islands | — | 0 | 0 | — | — | — | 0 | 0 | — | — |

C.N.M.I.: Commonwealth of Northern Mariana Islands.

U: Unavailable. —: No reported cases. N: Not notifiable. Cum: Cumulative year-to-date counts. Med: Median. Max: Maximum.

* Incidence data for reporting year 2006 is provisional.

† Updated weekly from reports to the Division of Vector-Borne Infectious Diseases, National Center for Zoonotic, Vector-Borne, and Enteric Diseases (proposed) (ArboNET Surveillance).

§ Contains data reported through the National Electronic Disease Surveillance System (NEDSS).

TABLE III. Deaths in 122 U.S. cities,* week ending November 25, 2006 (47th Week)

| Reporting Area | All causes, by age (years) | | | | | | | Reporting Area | All causes, by age (years) | | | | | | |
|-----------------------------|----------------------------|-------|-------|-------|------|----|------------------------|------------------------------|----------------------------|-------|-------|-------|------|-----|------------------------|
| | All Ages | ≥65 | 45-64 | 25-44 | 1-24 | <1 | P&I [†] Total | | All Ages | ≥65 | 45-64 | 25-44 | 1-24 | <1 | P&I [†] Total |
| New England | 458 | 314 | 104 | 23 | 10 | 7 | 38 | S. Atlantic | 669 | 405 | 163 | 57 | 25 | 19 | 39 |
| Boston, MA | 124 | 76 | 30 | 7 | 5 | 6 | 9 | Atlanta, GA | U | U | U | U | U | U | U |
| Bridgeport, CT | 24 | 18 | 6 | — | — | — | 5 | Baltimore, MD | 121 | 64 | 35 | 14 | 4 | 4 | 7 |
| Cambridge, MA | 15 | 13 | 2 | — | — | — | — | Charlotte, NC | 71 | 43 | 12 | 8 | 6 | 2 | 5 |
| Fall River, MA | 14 | 7 | 5 | 1 | 1 | — | 1 | Jacksonville, FL | 115 | 70 | 25 | 7 | 9 | 4 | 9 |
| Hartford, CT | 45 | 31 | 9 | 2 | 3 | — | 8 | Miami, FL | 53 | 31 | 13 | 6 | 3 | — | 1 |
| Lowell, MA | 17 | 12 | 3 | 2 | — | — | 1 | Norfolk, VA | 25 | 14 | 8 | — | 1 | 2 | 2 |
| Lynn, MA | 12 | 8 | 2 | 1 | 1 | — | 1 | Richmond, VA | 27 | 17 | 9 | 1 | — | — | 3 |
| New Bedford, MA | 24 | 19 | 4 | 1 | — | — | 1 | Savannah, GA | 31 | 19 | 7 | 3 | — | 2 | 2 |
| New Haven, CT | 43 | 30 | 11 | 1 | — | 1 | 3 | St. Petersburg, FL | 34 | 25 | 4 | 3 | — | 2 | 6 |
| Providence, RI | 48 | 36 | 8 | 4 | — | — | 3 | Tampa, FL | 106 | 73 | 20 | 10 | 2 | 1 | 3 |
| Somerville, MA | 3 | 2 | 1 | — | — | — | — | Washington, D.C. | 74 | 39 | 28 | 5 | — | 2 | — |
| Springfield, MA | 17 | 9 | 7 | 1 | — | — | 1 | Wilmington, DE | 12 | 10 | 2 | — | — | — | 1 |
| Waterbury, CT | 18 | 11 | 6 | 1 | — | — | 2 | E.S. Central | 742 | 487 | 170 | 57 | 13 | 15 | 54 |
| Worcester, MA | 54 | 42 | 10 | 2 | — | — | 3 | Birmingham, AL | 129 | 89 | 25 | 9 | 3 | 3 | 9 |
| Mid. Atlantic | 1,806 | 1,248 | 390 | 106 | 35 | 23 | 92 | Chattanooga, TN | 66 | 44 | 17 | 3 | — | 2 | 1 |
| Albany, NY | 37 | 18 | 12 | 5 | — | 2 | — | Knoxville, TN | 82 | 56 | 18 | 6 | 2 | — | 2 |
| Allentown, PA | 29 | 24 | 5 | — | — | — | 2 | Lexington, KY | 40 | 29 | 4 | 5 | 1 | 1 | 5 |
| Buffalo, NY | 82 | 58 | 21 | 3 | — | — | 6 | Memphis, TN | 212 | 122 | 63 | 19 | 4 | 4 | 24 |
| Camden, NJ | 28 | 14 | 11 | 2 | 1 | — | — | Mobile, AL | 82 | 60 | 16 | 4 | 1 | 1 | 3 |
| Elizabeth, NJ | 12 | 5 | 6 | — | — | 1 | 1 | Montgomery, AL | 31 | 17 | 10 | 2 | — | 2 | 5 |
| Erie, PA | 40 | 31 | 8 | — | 1 | — | 1 | Nashville, TN | 100 | 70 | 17 | 9 | 2 | 2 | 5 |
| Jersey City, NJ | 22 | 16 | 1 | 2 | 3 | — | 2 | W.S. Central | 924 | 559 | 235 | 74 | 27 | 29 | 41 |
| New York City, NY | 824 | 569 | 171 | 59 | 13 | 8 | 35 | Austin, TX | 58 | 32 | 14 | 9 | 2 | 1 | 5 |
| Newark, NJ | 33 | 16 | 8 | 3 | 4 | 2 | 1 | Baton Rouge, LA | 48 | 29 | 14 | 2 | 3 | — | — |
| Paterson, NJ | 18 | 9 | 8 | — | 1 | — | — | Corpus Christi, TX | 34 | 28 | 3 | 1 | 2 | — | 3 |
| Philadelphia, PA | 350 | 228 | 84 | 22 | 8 | 8 | 17 | Dallas, TX | 115 | 63 | 29 | 7 | 5 | 11 | 7 |
| Pittsburgh, PA [‡] | 19 | 13 | 4 | 2 | — | — | — | El Paso, TX | 62 | 41 | 12 | 6 | 2 | 1 | 5 |
| Reading, PA | 24 | 19 | 4 | — | 1 | — | — | Fort Worth, TX | 74 | 52 | 12 | 7 | — | 3 | 3 |
| Rochester, NY | 110 | 88 | 18 | 2 | 1 | 1 | 8 | Houston, TX | 206 | 113 | 64 | 16 | 6 | 7 | 5 |
| Schenectady, NY | 17 | 13 | 4 | — | — | — | 2 | Little Rock, AR | 50 | 27 | 14 | 7 | 1 | 1 | 2 |
| Scranton, PA | 23 | 19 | 3 | 1 | — | — | 2 | New Orleans, LA [¶] | U | U | U | U | U | U | U |
| Syracuse, NY | 86 | 72 | 11 | 2 | 1 | — | 13 | San Antonio, TX | 177 | 110 | 48 | 15 | 2 | 2 | 5 |
| Trenton, NJ | 21 | 12 | 6 | 1 | 1 | 1 | 1 | Shreveport, LA | 30 | 18 | 9 | 2 | 1 | — | 2 |
| Utica, NY | 16 | 12 | 3 | 1 | — | — | 1 | Tulsa, OK | 70 | 46 | 16 | 2 | 3 | 3 | 4 |
| Yonkers, NY | 15 | 12 | 2 | 1 | — | — | — | Mountain | 841 | 571 | 176 | 57 | 16 | 21 | 56 |
| E.N. Central | 1,579 | 1,060 | 346 | 111 | 33 | 29 | 95 | Albuquerque, NM | 88 | 55 | 22 | 8 | 2 | 1 | 8 |
| Akron, OH | U | U | U | U | U | U | U | Boise, ID | 30 | 21 | 7 | 2 | — | — | 5 |
| Canton, OH | 36 | 26 | 7 | 3 | — | — | 2 | Colorado Springs, CO | 41 | 29 | 5 | 3 | 1 | 3 | 4 |
| Chicago, IL | 257 | 127 | 74 | 40 | 12 | 4 | 24 | Denver, CO | 62 | 40 | 15 | 2 | 4 | 1 | 4 |
| Cincinnati, OH | 41 | 28 | 9 | 1 | 1 | 2 | 3 | Las Vegas, NV | 228 | 145 | 58 | 19 | 3 | 3 | 17 |
| Cleveland, OH | 203 | 143 | 47 | 9 | 3 | 1 | 13 | Ogden, UT | 19 | 13 | 1 | 2 | — | 3 | — |
| Columbus, OH | 153 | 104 | 29 | 11 | 3 | 6 | 10 | Phoenix, AZ | 140 | 94 | 24 | 13 | 2 | 7 | 7 |
| Dayton, OH | 94 | 71 | 18 | 2 | 3 | — | 4 | Pueblo, CO | 29 | 19 | 9 | 1 | — | — | 1 |
| Detroit, MI | 79 | 38 | 32 | 7 | 1 | 1 | 1 | Salt Lake City, UT | 103 | 75 | 20 | 5 | 2 | 1 | 3 |
| Evansville, IN | 27 | 25 | 1 | 1 | — | — | 2 | Tucson, AZ | 101 | 80 | 15 | 2 | 2 | 2 | 7 |
| Fort Wayne, IN | 44 | 26 | 12 | 2 | 3 | 1 | 2 | Pacific | 1,270 | 846 | 281 | 85 | 32 | 26 | 107 |
| Gary, IN | 11 | 6 | 3 | 2 | — | — | — | Berkeley, CA | 8 | 5 | 2 | — | 1 | — | 2 |
| Grand Rapids, MI | 64 | 45 | 11 | 3 | 1 | 4 | 10 | Fresno, CA | 87 | 52 | 23 | 8 | 3 | 1 | 8 |
| Indianapolis, IN | 187 | 122 | 40 | 15 | 5 | 5 | 10 | Glendale, CA | 6 | 1 | 4 | 1 | — | — | — |
| Lansing, MI | 33 | 26 | 5 | 2 | — | — | 3 | Honolulu, HI | 68 | 46 | 13 | 3 | — | 6 | 5 |
| Milwaukee, WI | 57 | 43 | 8 | 2 | — | 4 | 2 | Long Beach, CA | 60 | 39 | 14 | 1 | 5 | 1 | 2 |
| Peoria, IL | 45 | 32 | 9 | 3 | 1 | — | 3 | Los Angeles, CA | 110 | 46 | 36 | 21 | 5 | 2 | 3 |
| Rockford, IL | 79 | 54 | 21 | 3 | — | 1 | 3 | Pasadena, CA | 20 | 14 | 4 | — | 1 | 1 | 3 |
| South Bend, IN | 33 | 28 | 3 | 2 | — | — | 1 | Portland, OR | 100 | 69 | 23 | 4 | 4 | — | 12 |
| Toledo, OH | 93 | 76 | 16 | 1 | — | — | 1 | Sacramento, CA | 139 | 97 | 28 | 8 | 2 | 4 | 12 |
| Youngstown, OH | 43 | 40 | 1 | 2 | — | — | 1 | San Diego, CA | 84 | 54 | 21 | 6 | 3 | — | 9 |
| W.N. Central | 365 | 231 | 89 | 22 | 6 | 17 | 22 | San Francisco, CA | 146 | 95 | 41 | 8 | — | 2 | 20 |
| Des Moines, IA | 50 | 38 | 9 | 3 | — | — | 8 | San Jose, CA | 200 | 157 | 27 | 9 | 2 | 5 | 14 |
| Duluth, MN | 20 | 14 | 5 | 1 | — | — | 1 | Santa Cruz, CA | 25 | 21 | 4 | — | — | — | 2 |
| Kansas City, KS | 14 | 5 | 4 | 4 | — | 1 | — | Seattle, WA | 72 | 44 | 18 | 6 | 2 | 2 | 4 |
| Kansas City, MO | 67 | 45 | 11 | 3 | 2 | 6 | 1 | Spokane, WA | 47 | 35 | 7 | 3 | — | 2 | 7 |
| Lincoln, NE | 14 | 11 | 2 | 1 | — | — | 1 | Tacoma, WA | 98 | 71 | 16 | 7 | 4 | — | 4 |
| Minneapolis, MN | 32 | 20 | 9 | 2 | — | 1 | 1 | Total | 8,654** | 5,721 | 1,954 | 592 | 197 | 186 | 544 |
| Omaha, NE | 53 | 32 | 18 | 1 | — | 2 | 2 | | | | | | | | |
| St. Louis, MO | 58 | 29 | 18 | 5 | 4 | 2 | 6 | | | | | | | | |
| St. Paul, MN | 26 | 17 | 7 | — | — | 2 | — | | | | | | | | |
| Wichita, KS | 31 | 20 | 6 | 2 | — | 3 | 2 | | | | | | | | |

U: Unavailable. —: No reported cases.

* Mortality data in this table are voluntarily reported from 122 cities in the United States, most of which have populations of ≥100,000. A death is reported by the place of its occurrence and by the week that the death certificate was filed. Fetal deaths are not included.


† Pneumonia and influenza.

‡ Because of changes in reporting methods in this Pennsylvania city, these numbers are partial counts for the current week. Complete counts will be available in 4 to 6 weeks.

¶ Because of Hurricane Katrina, weekly reporting of deaths has been temporarily disrupted.

** Total includes unknown ages.

FIGURE I. Selected notifiable disease reports, United States, comparison of provisional 4-week totals November 25, 2006, with historical data



* No measles or rubella cases were reported for the current 4-week period yielding a ratio for week 47 of zero (0).
 † Ratio of current 4-week total to mean of 15 4-week totals (from previous, comparable, and subsequent 4-week periods for the past 5 years). The point where the hatched area begins is based on the mean and two standard deviations of these 4-week totals.

Notifiable Disease Data Team and 122 Cities Mortality Data Team
 Patsy A. Hall
 Deborah A. Adams Rosaline Dhara
 Willie J. Anderson Vernitta Love
 Lenee Blanton Pearl C. Sharp

The *Morbidity and Mortality Weekly Report (MMWR)* Series is prepared by the Centers for Disease Control and Prevention (CDC) and is available free of charge in electronic format. To receive an electronic copy each week, send an e-mail message to listserv@listserv.cdc.gov. The body content should read *SUBscribe mmwr-toc*. Electronic copy also is available from CDC's Internet server at <http://www.cdc.gov/mmwr> or from CDC's file transfer protocol server at <ftp://ftp.cdc.gov/pub/publications/mmwr>. Paper copy subscriptions are available through the Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; telephone 202-512-1800.

Data in the weekly *MMWR* are provisional, based on weekly reports to CDC by state health departments. The reporting week concludes at close of business on Friday; compiled data on a national basis are officially released to the public on the following Friday. Data are compiled in the National Center for Public Health Informatics, Division of Integrated Surveillance Systems and Services. Address all inquiries about the *MMWR* Series, including material to be considered for publication, to Editor, *MMWR* Series, Mailstop E-90, CDC, 1600 Clifton Rd., N.E., Atlanta, GA 30333 or to www.mmwrq@cdc.gov.

All material in the *MMWR* Series is in the public domain and may be used and reprinted without permission; citation as to source, however, is appreciated.

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services.

References to non-CDC sites on the Internet are provided as a service to *MMWR* readers and do not constitute or imply endorsement of these organizations or their programs by CDC or the U.S. Department of Health and Human Services. CDC is not responsible for the content of these sites. URL addresses listed in *MMWR* were current as of the date of publication.